

# RESTRICTED USE PESTICIDE

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

# QUALI-PRO

# Chlorpyrifos 4E

## *Insecticide*

**For control of listed insects infesting certain field, fruit, nut, and vegetable crops.**

ACTIVE INGREDIENT	% BY WT.
Chlorpyrifos: 0,0-diethyl-0-(3,5,6-trichloro-2-pyridinyl)phosphorothioate .....	42.5%
<b>INERT INGREDIENTS*:</b> .....	<u>57.5%</u>
<b>TOTAL:</b> .....	100.0%

Contains 4 pounds of chlorpyrifos per gallon.

\*Contains petroleum distillates.

Contains chlorpyrifos, the active ingredient used in Lorsban® 4E.

Quali-Pro Chlorpyrifos 4E is not manufactured or distributed by Dow AgroSciences LLC.

EPA Reg. No. 66222-19

**KEEP OUT OF REACH OF CHILDREN**  
**WARNING-AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### WARNING

May be fatal if swallowed. Causes substantial but temporary eye injury. Causes skin irritation. Harmful if absorbed through skin. Do not get in eyes, on skin, or on clothing.

For additional precautionary, handling, and use statements, see inside of this booklet.

Manufactured for:  
**Makhteshim Agan of North America, Inc.**  
3120 Highwoods Blvd, Suite 100  
Raleigh, NC 27604

EPA 112509/Notif 031710/102111/Rev D

**FIRST AID  
ORGANOPHOSPHATE**

<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Do not give any liquid to a person.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For medical emergencies involving this product, call Prosar at 1-877-250-9291.

**NOTE TO PHYSICIAN:** This product contains an organophosphate that inhibits cholinesterase. Treat symptomatically. If exposed, plasma and red blood cell cholinesterase tests may indicate significance of exposure (baseline data are useful). Atropine, only by injection, is the preferable antidote. Oximes, such as 2-PAM/protopam, may be therapeutic if used early; however, use only in conjunction with atropine. In case of severe acute poisoning, use antidote immediately after establishing an open airway and respiration. Contains petroleum distillate. Vomiting may induce aspiration pneumonia.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
WARNING**

May be fatal if swallowed. Causes substantial but temporary eye injury. Causes skin irritation. Harmful if absorbed through skin. Do not get in eyes, on skin, or on clothing.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Some materials that are chemical resistant to this product are listed below. If you want more options, follow the instructions for category *H* on an EPA chemical resistance category selection chart.

**All mixers, loaders, other applicators, and other handlers must wear:**

- Coveralls over long-sleeved shirt and long pants
- Chemical-resistant gloves
- Chemical-resistant apron when mixing or loading or exposed to the concentrate
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure
- A NIOSH-approved dust mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any R, P, or HE filter

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

## ENGINEERING CONTROLS

When handlers use closed systems or closed cab motorized ground application equipment in a manner that meets the requirements listed in Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Remove and wash contaminated clothing before reuse.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, small mammals, and birds. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment washwater or rinsate.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. Protective information may be obtained from your cooperative agricultural extension service.

### PHYSICAL OR CHEMICAL HAZARDS

Combustible. Do not use or store near heat or open flame.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

Read all **DIRECTIONS FOR USE** carefully before applying.

### AGRICULTURAL USE REQUIREMENTS

Use this product in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of:

- 4 days for fruit trees
- 5 days for citrus
- 3 days for cauliflower
- 24 hours for all other crops not listed above

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated such as plants, soil, or water is:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant gloves made out of any waterproof material
- Chemical-resistant footwear plus socks
- Protective eyewear
- Chemical-resistant headgear for overhead exposure

Certified crop advisors or persons entering under their direct supervision under certain circumstances may be exempt from the early reentry requirement pursuant to 40 CFR Part 170.

Notify workers of the application by warning them orally and by posting warning signs at entrances to treated areas.

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides [40 CFR Part 170]. The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep children, pets, and other unprotected persons out of treated area until sprays have dried.

### USE INFORMATION

Quali-Pro Chlorpyrifos 4E insecticide forms an emulsion when diluted with water and is suitable for use in all conventional spray equipment. Consult your State Experiment Station or State Extension Service for proper timing of applications.

When an adjuvant is to be used with this product, Makhteshim Agan of North America, Inc. suggests the use of a Chemical Producers and Distributors Association certified adjuvant.

### USE PRECAUTIONS

Do not formulate this product into other end use products. **Attention:** Do not cut or weld container.

### SPRAY DRIFT

Do not allow spray to drift from the application site and contact people, structures people occupy at any time and the associated property, parks and recreation areas, non-target crops, aquatic and wetland areas, woodlands, pastures, rangelands, or animals.

- For ground boom applications, do not apply within 25 feet of rivers, natural ponds, lakes, streams, reservoirs, marshes, estuaries, and commercial fish ponds. Apply with nozzle height no more than 4 feet above the ground or crop canopy and when wind speed is 10 mph or less at the application site, as measured by an anemometer. Use fine or coarser spray according to ASAE 572 definition for standard nozzles or VMD for spinning atomizer nozzles.
- For orchard/vineyard airblast applications, do not apply within 50 feet of rivers, natural ponds, lakes, streams, reservoirs, marshes, estuaries,

and commercial fish ponds. Direct spray above trees/vines and turn off outward-pointing nozzles at row ends and outer rows. Apply only when wind speed is 3 to 10 mph at the application site, as measured by an anemometer outside of the orchard/vineyard on the upwind side.

- For aerial applications, do not apply within 150 feet of rivers, natural ponds, lakes, streams, reservoirs, marshes, estuaries, and commercial fish ponds. The boom width must not exceed 75% of the wingspan or 90% of the rotary blade. Use upwind swath displacement and apply only when wind speed is 3 to 10 mph, as measured by an anemometer. Use fine or coarser spray according to ASAE 572 definition for standard nozzles or VMD for spinning atomizer nozzles. If application includes a no-spray zone, do not release spray at a height greater than 10 feet above the ground or the crop canopy.
- For overhead chemigation, do not apply within 25 feet of rivers, natural ponds, lakes, streams, reservoirs, marshes, estuaries, and commercial fish ponds. Apply only when wind speed is 10 mph or less.

The applicator also must use all other measures necessary to control drift.

### SPRAY MIX DIRECTIONS

To prepare the spray, add a portion of the required amount of water to the spray tank and with agitation, add the Quali-Pro Chlorpyrifos 4E. Complete filling the tank with the balance of water needed. Maintain sufficient agitation during both mixing and application to ensure uniformity of the spray mixture.

Quali-Pro Chlorpyrifos 4E may also be used in tank mixtures with certain herbicides and/or with non-pressure fertilizer solutions as recommended under specific crop use directions. Prepare tank mixtures in the same manner as recommended above for use of Quali-Pro Chlorpyrifos 4E alone. When tank mixtures of Quali-Pro Chlorpyrifos 4E and herbicides are involved, add wettable powders first, flowables second, and emulsifiable concentrates last. Where a fertilizer solution is involved, it is strongly recommended that a fertilizer pesticide compatibility agent such as Unite or Complex be used. Maintain constant agitation during both mixing and application to ensure uniformity of the spray mixture. Do not allow spray mixtures to stand overnight.

**NOTE:** Test compatibility of the intended tank mixture before adding Quali-Pro Chlorpyrifos 4E to the spray or tank mix. Add proportionate amounts of each ingredient to a pint or quart jar, cap, shake, and let set 15 minutes. Formation of precipitates that do not readily redisperse indicates an incompatible mixture that should not be used.

### SPRINKLER IRRIGATION

Quali-Pro Chlorpyrifos 4E may be applied by sprinkler irrigation for the following crop uses: alfalfa, almond (orchard floors only), citrus orchard floors, corn (field and sweet), cotton, cranberry, mint (peppermint and spearmint), pecan and walnut orchard floors, sorghum, soybeans, sugar beet, and wheat.

See the use sections for the individual crops for further application information. Do not apply this product to the above-listed crops through any other type of irrigation system. Do not apply this product by chemigation to any other crop.

### SPRINKLER USE DIRECTIONS

The following use directions are to be followed when Quali-Pro Chlorpyrifos 4E is applied through sprinkler irrigation systems. Thoroughly clean the injection system and tank of any fertilizer or chemical residues, and dispose of the residues according to State and Federal laws. Flush the injector with soap and water. Determine the amount of insecticide needed to cover the desired acreage. Pump the required Quali-Pro Chlorpyrifos 4E into a steel tank, start mechanical or hydraulic agitation, and add in order the non-emulsifiable oil and/or water. Continually agitate the mixture containing Quali-Pro Chlorpyrifos 4E. Set the sprinkler system to deliver the desired inches of water per acre. Start the water pump and sprinkler, and let the system achieve the desired pressure and speed before starting the injector. Start the injector and calibrate the injector system according to number 14 in **SPECIAL USE PRECAUTIONS** section. The mixture containing Quali-Pro Chlorpyrifos 4E must be injected continuously and uniformly into the irrigation water line as the sprinkler is moving. This procedure is necessary to deliver the desired rate per acre in a uniform manner. When the application is finished, allow the entire irrigation and injector system to be thoroughly flushed clean before stopping the system.

### SPRINKLER USE PRECAUTIONS

The following use precautions will result in a safe and successful application of mixture containing Quali-Pro Chlorpyrifos 4E.

1. Apply this product only through sprinkler irrigation systems including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, micro sprinkler, or hand move. Do not apply this product through any other type of irrigation system.

2. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.
3. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers, or other experts.
4. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system.
5. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.
6. The system must contain a functional check valve, vacuum-relief valve, and a low-pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. Refer to the American Society of Agricultural Engineer's Engineering Practice 409 for more information.
7. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
8. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
9. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
10. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
11. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. The metering pump must provide a greater pressure than that of the irrigation system at the point of injection. The pump must meet Section 675 for "Electrically Driven or Controlled Irrigation Machines" NEC 70 and must contain Viton or Teflon seals.
12. To insure uniform mixing of the insecticide into the water line, inject the mixture through a nozzle placed in the fertilizer injection port or just ahead of an elbow or tee in the irrigation line so that the turbulence created at those points will assist in mixing. It is suggested that the injection point be higher than the insecticide tank to prevent siphoning.
13. The steel tank holding the insecticide mixture should be large enough to allow the system to complete a revolution with one filling. It should be free of rust, fertilizer sediment, and foreign material, and equipped with an in-line strainer situated between the tank and the injector pump.
14. In order to calibrate the irrigation system and injector to apply the mixture containing Quali-Pro Chlorpyrifos 4E, determine the following: 1) Calculate the number of acres irrigated by the system; 2) Set the irrigation rate and determine the number of minutes for the system to cover the intended treatment area; 3) Calculate the total gallons of insecticide mixture needed to cover the desired acreage. Divide the total gallons of insecticide mixture needed by the number of minutes to cover the treatment area. This value equals the gallons per minute output that the injector must deliver. Convert the gallons per minute to milliliters or ounces per minute. Calibrate the injector pump with the system in operation at the desired irrigation rate. It is suggested that the injector pump be calibrated at least twice before operation, and monitor the system.
15. Do not apply when wind speed favors drift beyond the area intended for treatment. End guns must be turned off during the application if they irrigate non-target areas.
16. Do not allow irrigation water to collect or run off and pose a hazard to livestock, wells, or adjoining crops.
17. Allow foliage to dry before reentering the field.
18. Do not apply through sprinkler systems that deliver a low coefficient of uniformity such as certain water drive units.

## APPROVED CROPS

### ALFALFA

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E to control the following pests at the dosages indicated by application as a broadcast foliar spray:

PEST	QUALI-PRO CHLORPYRIFOS 4E
Aphids (suppression)	1/2 pint/acre (0.25 lb. a.i./acre)
Corn rootworm adults (spotted cucumber beetle), Grasshoppers, Leafhoppers	1/2 – 1 pint/acre (0.25 – 0.5 lb. a.i./acre)
Alfalfa blotch leafminer, Alfalfa looper, Alfalfa weevil larvae and adults, Armyworms, Cutworms, Egyptian alfalfa weevil larvae and adults, Plant bugs, Leafhoppers, Spittlebugs	1 – 2 pints/acre (0.5 – 1 lb. a.i./acre)

**NOTE:** Use higher rates to control spotted alfalfa aphid in California and Nevada. Stubble spray may be applied to control leafhopper in the Northeast.

Mix the required dosage with enough water to ensure thorough coverage of crop foliage and apply using aerial (fixed-wing or helicopter) or power-operated ground spray equipment. For aerial application, use 2 to 5 gallons of water per acre. For best coverage when using ground application, a minimum of 20 gallons of water per acre with hollow cone nozzles is recommended. Control may be reduced at low spray volumes under high temperature and wind conditions. Treat when field counts or crop injury indicates that damaging pest populations are developing or present; however, do not apply more than once per crop cutting. Some reduction in insect control may be evident under excessively cool conditions. For Egyptian alfalfa weevil control in California, apply the specified dosage in a minimum of 5 gallons of water per acre when larvae are actively feeding and populations reach 15 to 20 larvae per 180° sweep with a 15-inch diameter net.

Quali-Pro Chlorpyrifos 4E may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above-listed foliar pests. For best results, use the listed rate of Quali-Pro Chlorpyrifos 4E per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See **SPRINKLER IRRIGATION** section for further information.

Do not tank mix Quali-Pro Chlorpyrifos 4E with pesticides, surfactants, or fertilizer formulations unless prior use has shown the combination non-injurious under your current conditions of use. Some phytotoxic symptoms, may be observed on young, tender, rapidly growing alfalfa when treated with Quali-Pro Chlorpyrifos 4E. Alfalfa will outgrow the symptoms and no yield loss should be expected.

This product is highly toxic to bees exposed to direct treatment on alfalfa. Do not apply if nearby bees are clustered outside of hives and bees are foraging. Protective information may be obtained from your agricultural extension service.

**RESTRICTIONS:** Do not cut or graze treated alfalfa within 7 days after application of 1/2 pint (0.25 lb. a.i.) of Quali-Pro Chlorpyrifos 4E per acre, within 14 days after application of 1 pint (0.5 lb. a.i.) per acre, or within 21 days after application of rates above 1 pint (0.5 lb. a.i.) per acre. Do not make more than 4 applications per year or apply more than once per crop cutting. Do not make a second application of Quali-Pro Chlorpyrifos 4E within 10 days of the first application. To avoid contamination of irrigation floodwaters, do not flood irrigate within 24 hours following an application of Quali-Pro Chlorpyrifos 4E.

*The maximum single application rate is 1.0 lb. a.i. chlorpyrifos per acre.*

## APPLE TREE TRUNK

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 days unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E to control American plum borer, apple bark borer, broad necked root borer, dogwood borer, flatheaded apple-tree borer, and roundheaded appletree borer. Mix 1 1/2 qts. of Quali-Pro Chlorpyrifos 4E with 100 gallons of water; and using a low-volume handgun or shielded spray, apply directly to the trunk from a distance no further than 4 ft. Do not allow spray to contact foliage or fruit.

**RESTRICTIONS:** For use only in states East of the Rockies (except Mississippi). Do not apply within 28 days before harvest. Make no more than one application to the tree trunk per year as either a pre-bloom or post-bloom application. Do not apply if a pre-bloom application has been made during the year using any other chlorpyrifos product. Do not allow meat or dairy animals to graze in treated orchards. Treat the lower 4 feet of the trunk only. If wind speed is greater than 10 mph, do not apply.

## ASPARGUS

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E to control armyworms, asparagus aphids, asparagus beetles, cutworms, and grasshoppers by application at the rate of 2 pints per acre (1 lb. a.i.). Mix the specified dosage in sufficient water to ensure thorough coverage of treated plants and apply a broadcast foliar spray. For cutworms, it is preferable to apply Quali-Pro Chlorpyrifos 4E when the soil is moist and worms are active on or near the soil surface. Applications may be made during the fern stage for control of asparagus beetles and asparagus aphids when field counts or crop injury indicates that damaging pest populations are developing or present.

**RESTRICTIONS:** Do not make more than one preharvest application per season or apply within one day of harvest. Do not make more than two postharvest applications during the fern stage. Do not make a second application of Quali-Pro Chlorpyrifos 4E within 10 days of the first application. Based on available residue data, the use of Quali-Pro Chlorpyrifos 4E on asparagus is limited to the Midwest and Pacific Northwest (AZ, CA, ID, IL, IA, KS, MI, MN, MO, NE, ND, OH, OR, SD, WA, WI). The maximum single application rate preharvest or postharvest is 1.0 lb. a.i. chlorpyrifos per acre.

## CHERRIES (TRUNK SPRAY)

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 days unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E for the control of lesser peach tree borer, greater peach tree borer, and American plum borer by application as a trunk spray. Mix 1 1/2 to 3 quarts (1.5 to 3 lbs. a.i.) of Quali-Pro Chlorpyrifos 4E with 100 gallons of water and apply as a coarse, low-pressure spray to give uniform coverage of tree trunks and lower limbs. Make a second application two weeks after the first one and a third application after harvest. Avoid contact with foliage in sweet cherries, as premature leaf drop may result. Consult your State Agricultural Experiment Station or Extension Service specialist for proper time to treat in your area.

In addition, one of the three allowable applications per year may be applied as a dormant spray for control of San Jose scale, peach twig borer, and climbing cutworm. For control of these pests, tank mix 1/2 to 1 pint (0.25 to 0.5 lb. a.i.) of Quali-Pro Chlorpyrifos 4E with 1 to 2 gallons of a petroleum oil recommended for dormant use in 100 gallons of water and spray the entire tree by application to runoff using ground spray equipment. For low-volume (concentrate) sprays (40 to 100 gallons of spray mixture per acre), use the same amounts of Quali-Pro Chlorpyrifos 4E and spray oil per acre required for application as a dilute spray and apply in a manner that will ensure thorough coverage of the trees. Use the higher dosage of Quali-Pro Chlorpyrifos 4E for severe infestations. Use oil as recommended by your State Agricultural Experiment Station or Extension Service specialist.

**RESTRICTIONS:** Make only three applications per year. Do not apply within 21 days before harvest. Do not allow meat or dairy animals to graze in treated orchards.



## CHRISTMAS TREE PLANTATIONS

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E at the rate indicated to control the following insects on the tree varieties listed.

TREE VARIETY	PEST	DOSAGE QUALI-PRO CHLORPYRIFOS 4E	REMARKS
Balsam fir Concolor fir Douglas fir Eastern white pine Fraser fir Grand fir Noble fir Scotch pine White spruce	Ants (except fire ants, carpenter ants, harvester ants, and pharaoh ants), Aphids, Adelgids (Cooley, Eastern spruce gall), European pine shoot moth, European pine sawfly, Grasshopper, Gypsy moth, Mites (European red spider, Twospotted spider) [except in WA & OR], Pales weevil (adult), Pine needle midge, Pine spittlebug, Plant bugs, Spittlebugs, Spruce budworm, Spruce needleminer, Scale (Pine needle, Pine tortoise, Spruce bud, Black pine, Striped pine)	1 quart/acre (1 lb. a.i./acre)	Do not treat plants under extreme heat and drought stress.  Apply as a foliar spray using ground equipment. Thorough coverage of foliage is essential. Use a minimum 10 gpa of finished spray.  For effective control of adult spider mites if large numbers of eggs are present, apply a second spray 7 to 10 days after initial treatment to control newly hatched nymphs.  For scale control, apply when scale crawlers are active.  Before treating other conifer species, make application and observe for 7 to 10 days for symptoms of phytotoxicity. The user is responsible for determining if it is safe to treat other conifer species under commercial-growing conditions.
	Pales weevil	3 quarts/100 gals. (3 lbs. a.i./100 gals.)	Apply as a cut stump drench.

**RESTRICTIONS:** Use Quali-Pro Chlorpyrifos 4E on tree plantations only in Connecticut, Maine, Maryland, Massachusetts, Michigan, New Hampshire, New Jersey, New York, North Carolina, Ohio, Oregon, Pennsylvania, South Carolina, Tennessee, Vermont, Virginia, Washington, and Wisconsin. Do not allow livestock to graze in treated areas. Do not make more than 3 applications per season. Do not make a second application of Quali-Pro Chlorpyrifos 4E within 7 days of the first application. Do not apply by aerial application.

## CITRUS FRUITS

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 5 days unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E at the rates indicated according to the designated geographic area to control the following pests. Use the lower rates for light infestations and increase the dosage for heavier infestations. A petroleum spray oil recommended for use on citrus trees may be added to dilute spray mixtures only at a rate of up to 1.8 gallons per 100 gallons of water to improve control of aphids, mealybugs, scale insects, and thrips. Treat when insects become a problem or in accordance with the local spray schedule recommended by your state extension service specialist.

Quali-Pro Chlorpyrifos 4E may be applied in tank mixtures with ethion, dicofol, Agri-Mek, or Vendex. See **MIXING DIRECTIONS** for further instructions. Read and carefully follow all applicable directions, restrictions, and precautions on labeling for the other products used in combination with Quali-Pro Chlorpyrifos 4E.

**PRECAUTIONS:** Observe local use directions for tank mix combinations, especially in regard to applications of Quali-Pro Chlorpyrifos 4E plus

spray oil. Consult with a county farm advisor, county agency, extension service personnel, agricultural commissioner, or pest control advisor for such information regarding a given locality.

Do not apply when trees are stressed by drought or high temperatures.

Do not tank mix Quali-Pro Chlorpyrifos 4E with Difolatan 80 Sprills, as crop injury may occur.

Quali-Pro Chlorpyrifos 4E is highly toxic to bees exposed to direct treatment. Do not apply when bees are actively visiting the area. During the bloom period in California, apply from one hour after sunset until two hours before sunrise.

**RESTRICTIONS:** Do not apply more than 2 applications (does not include citrus orchard floor) or more than 15 pints (7.5 lbs. a.i.) of Quali-Pro Chlorpyrifos 4E per acre per year. Do not make second foliar application within 30 days of the first application. Do not treat within 21 days of harvest for applications up to 7 pints (3.5 lbs. a.i.) of Quali-Pro Chlorpyrifos 4E per acre nor within 35 days for application of rates above 7 pints (3.5 lbs. a.i.) per acre. Do not allow livestock to graze in treated areas. The use of application rates greater than 8 pints (4 lbs. a.i.) per acre is allowed only in the following California counties: Fresno, Tulare, Kern, Kings, and Madera. Do not allow meat or dairy animals to graze in treated areas.

CROP	GEO-GRAPHIC LOCATION	PEST	DOSAGE PT./ACRE	SPRAY VOLUME GAL./ACRE	REMARKS
Calamondin, Chironja, Citrus Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemons, Limes, Mandarin (tangerine), Oranges, Pummelo, Satsuma mandarin, Tangelo, Tangor, and Other Citrus Fruit	California – Fresno, Tulare, Kern, Kings, and Madera Counties Only	Red scale	8 – 12 (4 – 6 lbs. a.i.)	Ground: 100 – 2400	Do not use a spray concentration of Quali-Pro Chlorpyrifos 4E of less than 1/2 pt./100 gals. of total volume. <b>Additional Precautions for California and Arizona:</b> Do not use Quali-Pro Chlorpyrifos 4E in combination with spray oil when temperatures are expected to exceed 95°F the day of application or for several consecutive days thereafter. Do not apply during the months of December, January, or February.
Calamondin, Chironja, Citrus Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemons, Limes, Mandarin (tangerine), Oranges, Pummelo, Satsuma mandarin, Tangelo, Tangor, and Other Citrus Fruit	California, Arizona	Aphids, Katydid, Lepidopterous larvae, Avocado leafroller, Cutworms, Fruit tree leafroller, Orange tortrix, Western tussock moth	2 – 7 (1 – 3.5 lbs. a.i.)	Ground: 100 – 750  Aerial: min. 15	
		Scale insects (Black scale, Brown soft scale, California red scale)	8 (4 lbs. a.i.)	100 – 2400	
		Thrips (suppression), Mealybugs	6 – 12 (3 – 6 lbs. a.i.)	100 – 750	

(continued on next page)

**CITRUS FRUITS** (continued)

<b>CROP</b>	<b>GEO-GRAPHIC LOCATION</b>	<b>PEST</b>	<b>DOSAGE PT./ACRE</b>	<b>SPRAY VOLUME GAL./ACRE</b>	<b>REMARKS</b>
Calamondin, Chironja, Citrus Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemons, Limes, Mandarin (tangerine), Oranges, Pummelo, Satsuma mandarin, Tangelo, Tangor, and Other Citrus Fruit	Florida	Aphids, Grasshoppers, Orangedogs, Mealybugs, Scale insects (Snow scale, Florida red scale, Purple scale, Long scale, Chaff scale, Black scale, Brown soft scale)	2 - 4 (1 - 2 lbs. a.i.)	Ground: 100 - 1400  Aerial: min. 20	Do not use a spray concentrate of Quali-Pro Chlorpyrifos 4E of less than 1/2 pt./100 gals. of water per acre. Lubber grasshoppers must be controlled when they are small (less than 1 inch in length) by direct contact with spray.
		Citrus rust mites	4 - 7 (2 - 3.5 lbs. a.i.)	100 - 700	Do not use a spray concentration of Quali-Pro Chlorpyrifos 4E of less than 1 pt./100 gals. of water per acre.
Calamondin, Chironja, Citrus Citron, Citrus Hybrids, Grapefruit, Kumquat, Lemons, Limes, Mandarin (tangerine), Oranges, Pummelo, Satsuma mandarin, Tangelo, Tangor, and Other Citrus Fruit	Texas	Aphids, Cutworms, Katydid, Mealybugs, Scale insects (Brown soft scale, California red scale, Chaff scale)	4 - 7 (2 - 3.5 lbs. a.i.)	200 - 700	Do not use less than 1/2 pt. of Quali-Pro Chlorpyrifos 4E per 100 gals. of water in dilute applications.
		Citrus rust mites (suppression)	4 - 7 (2 - 3.5 lbs. a.i.)	200 - 700	
Small Transplanted Grapefruit, Orange, and Other Citrus Trees	Texas	Aphids, Cutworms, Katydid, Mealybugs, Scale insects (Brown soft scale, California red scale, Chaff scale)	Max of 7 (3.5 lbs. a.i.)		Apply Quali-Pro Chlorpyrifos 4E at a rate of 1 fl. oz./1 gal. of water with a backpack sprayer. Apply to runoff.

## CITRUS ORCHARD FLOORS

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 5 days unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E to control ant species (except fire ants, carpenter ants, harvester ants, and pharaoh ants) by applying the specified dose in 25 or more gallons of water with ground application equipment that will uniformly apply the spray to the orchard floor. Do not allow spray to contact foliage or fruit. To control foraging ants, apply Quali-Pro Chlorpyrifos 4E to the orchard floor at the rate of 3/4 to 1 quart (0.75 to 1 lb. a.i.) per acre. Repeat as needed. For best insect control, uniform coverage of the orchard floor is necessary. Do not apply where weed growth or other obstructions would impede uniform coverage of the orchard floor. Do not apply in tank mixtures with Evik herbicide. Foliar applications of Quali-Pro Chlorpyrifos 4E may be made in addition to the orchard floor, but must comply with the 10-day re-treatment interval.

Quali-Pro Chlorpyrifos 4E may also be applied to citrus orchard floors through sprinkler irrigation systems only if the system uniformly covers the soil surface at the base of the tree. Use the listed amount of Quali-Pro Chlorpyrifos 4E per acre. See **SPRINKLER IRRIGATION** section for further information.

**RESTRICTIONS:** Do not apply last treatment within 28 days before harvest. Do not allow livestock or dairy animals to graze in treated areas. Do not apply more than 3 quarts (3 lbs. a.i.) per acre per year. Do not make more than 3 applications per year (does not include foliar applications to citrus trees). Do not make a second application within 10 days of the first application. The maximum single application rate is 1 lb. a.i. chlorpyrifos per acre.

## CORN (FIELD), CORN (SWEET) (INCLUDING CORN GROWN FOR SEED)

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

### Conservation Tillage: Preplant, At-Plant, or Preemergence Applications

Apply as a broadcast spray to surface trash and exposed soil using power-operated ground spray equipment. Use a total spray volume of 20 gpa or more. Use a higher rate in the rate range to extend residual control.

**Tank Mixing:** Quali-Pro Chlorpyrifos 4E may also be applied in tank mixtures with paraquat or glyphosate and/or liquid fertilizer solutions. See **MIXING DIRECTIONS** section in this label for tank mixing instructions. Read and carefully follow all applicable directions, restrictions, and precautions on labeling for each product used in combination with Quali-Pro Chlorpyrifos 4E.

TARGET PESTS	QUALI-PRO CHLORPYRIFOS 4E
Armyworms, Cutworms	1 – 2 pints/acre (0.5 – 1 lb. a.i./acre)

### Postemergence Application

Apply as a postemergence broadcast spray using sufficient spray volume to ensure thorough coverage of treated plants, but no less than 15 gallons per acre (gpa) for ground spray equipment or 2 to 5 gpa for aircraft equipment. Control may be reduced at low spray volumes under high temperature and wind conditions. Quali-Pro Chlorpyrifos 4E may be tank mixed with glyphosate products such as Glyphomax® XRT herbicide or Durango™ herbicide when application is to be made to glyphosate-tolerant corn. **NOTE:** Do not apply aerially in Mississippi.

**Chemigation:** Quali-Pro Chlorpyrifos 4E may be broadcast applied postemergence through sprinkler irrigation systems at listed application rates to control listed foliar pests. For best results, tank mix Quali-Pro Chlorpyrifos 4E with 2 pints of non-emulsifiable oil. See **SPRINKLER IRRIGATION** section of this product label for application instructions.

TARGET PESTS	QUALI-PRO CHLORPYRIFOS 4E
Grasshoppers	1/2 – 1 pint/acre (0.25 lb. – 0.5 lb. a.i./acre)
Aphids, Armyworms, Chinch bugs (1), Corn rootworm adults (2), Cutworms (3), European corn borer (5), Flea beetle adults (1), Southern corn leaf beetle, Webworms (4), Western bean cutworm	1 – 2 pints/acre (0.5 – 1 lb. a.i./acre)
Corn earworm, Southwestern corn borer (6)	1 1/2 – 2 pints/acre (0.75 – 1 lb. a.i./acre)
Billbugs (1), Common stalk borer (9), Corn rootworm larvae (7),(8), Lesser cornstalk borer	2 pints/acre (1 lb. a.i./acre)

#### **Pest Specific Use Directions:**

- For best billbug, chinch bug, or flea beetle control, ground apply in a minimum spray volume of 20 to 40 gpa at 40 psi. If corn is less than 6 inches tall, apply in a 9- to 12-inch wide band over the row. For corn greater than 6 inches tall, apply using drop nozzles directed to the base of the plant. Do not reduce the application rate for banded or directed applications. Concentrate the full labeled dosage rate in the treated zone. When chinch bugs continue to immigrate to corn over a prolonged period or under extreme pest pressure, a second application may be needed.
- The listed dosage will control silk clipping by corn rootworm adults.
- For cutworms, it is preferable to apply Quali-Pro Chlorpyrifos 4E when soil is moist and worms are active on or near the soil surface. If ground is dry, cloddy, or crusted at time of treatment, worms may be protected from the spray and effectiveness will be reduced. Shallow incorporation using a rotary hoe or other suitable equipment immediately before or soon after treatment may improve control. A second application may be required if damage or density levels exceed economic thresholds established for your area.
- For webworm control, shallow incorporation using a rotary hoe or other suitable equipment immediately before or soon after treatment is necessary.
- For European corn borer control, use 1 1/2 to 2 pints per acre when application is made with power-operated ground or aerial equipment or 1 to 2 pints per acre when application is made through a sprinkler irrigation system. University research indicates that achieving greater than 50% control of first-generation European borer with a single liquid insecticide treatment is highly dependent upon timing, insecticide placement, and weather conditions.
- For southwestern corn borer, a second application may be applied 21 days later if needed due to reinfestation.
- For postemergence control of corn rootworm larvae, apply at cultivation. Direct the spray to both sides of the row at the base of the plants just ahead of the cultivator shovels. Cover the insecticide with soil around the brace roots. A cultivation application of Quali-Pro Chlorpyrifos 4E may be made in addition to an at-planting application of Chlorpyrifos 15G.
- Quali-Pro Chlorpyrifos 4E may also be applied through sprinkler irrigation systems at the rate of 2 pints per acre to control corn rootworm larvae. Time the application to coincide with the appearance of the second instar larvae. Apply with enough water to wet the root zone to the depth control needed. If soils are wet, allow enough soil drying to occur such that an application using a minimum amount of water will not produce surface runoff. See Chemigation (**SPRINKLER IRRIGATION**) section of this label for application instructions.
- Do not use Quali-Pro Chlorpyrifos 4E in combination with a burndown herbicide for control of common stalk borer. For common stalk borer control, treat approximately 11 days after application of glyphosate or after burndown with paraquat herbicide is complete (3 to 5 days).

#### **RESTRICTIONS – CORN (FIELD), CORN (SWEET) (INCLUDING CORN GROWN FOR SEED):**

Do not apply within 21 days before harvest of grain or ears. Do not apply more than 3 lbs. a.i. chlorpyrifos (6 pints of Quali-Pro Chlorpyrifos 4E) per acre per season. Do not make more than 3 applications per season of any product containing chlorpyrifos including the maximum allowed of 2 granular applications at the 1 lb. a.i. chlorpyrifos rate. Do not make a second application of Quali-Pro Chlorpyrifos 4E or other product containing chlorpyrifos within 10 days of the first application. The maximum single application rate is 1 lb. a.i. chlorpyrifos (2 pints of Quali-Pro Chlorpyrifos 4E) per acre. Do not apply in tank mixes with Steadfast or Lightning herbicides. If more than 1 lb. a.i. granular chlorpyrifos per acre is applied at-plant (for a maximum of 1.3 lbs. a.i. per acre per season), only 1 additional application of a liquid product containing chlorpyrifos at 1 lb. a.i. per acre is allowed per season, for a total of 2.3 lbs. a.i. chlorpyrifos per acre per season.

## COTTON

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E for control of the following pests in all states except Arizona and California at the dosages indicated:

PEST	QUALI-PRO CHLORPYRIFOS 4E
Cotton fleahopper, Plant bugs (Lygus, Mirids)	3/8 – 1 pint/acre (0.18 – 0.5 lb. a.i./acre)
Fall armyworm, Grasshopper, Thrips, Yellowstriped armyworm	1/2 – 1 pint/acre (0.25 – 0.5 lb. a.i./acre)
Cotton aphid	1/2 – 2 pints/acre (0.25 – 1 lb. a.i./acre)
Spider mites	1 pint/acre (0.5 lb. a.i./acre)
Beet armyworm, Cotton budworm, Tobacco budworm, Cutworms, Pink bollworm, Saltmarsh caterpillar	1 1/2 – 2 pints/acre (0.75 – 1 lb. a.i./acre)

**NOTE:** The listed dosage rate of 3/8 pint per acre will not achieve the high degree of control of the highest label rate but will minimize the damage done by plant bugs and cotton fleahopper and allow the beneficial insects to survive, build up, and be available to aid in the control of bollworms infesting cotton. For infestations of cotton aphids that are difficult to control, use a higher dosage within the indicated rate range.

Use Quali-Pro Chlorpyrifos 4E for control of the following pests in Arizona and California at the dosages indicated:

PEST	QUALI-PRO CHLORPYRIFOS 4E
Armyworms, Cotton aphid, Cotton fleahopper, Lygus, Saltmarsh caterpillar, Thrips	1 – 2 pints/acre (0.5 – 1 lb. a.i./acre)
Cotton bollworm, Tobacco budworm, Boll weevil, Cutworms, Pink bollworm	2 pints/acre (1 lb. a.i./acre)

**NOTE:** The 2-pint rate will aid in the suppression of cotton leafperforator and spider mites.

Mix the required dosage with sufficient water to ensure thorough coverage of plants and apply using aerial or power-operated ground spray equipment. For aerial application, use at least 2 gallons of spray per acre. For ground application, use sufficient spray volume to ensure thorough coverage of treated plants but not less than 10 gallons of spray per acre. Increase spray volume when foliage is dense and/or when pest populations are high, and/or under high temperature and wind conditions. Treat when field counts indicate damaging insect populations are developing or present. Re-treat as necessary to maintain control.

Quali-Pro Chlorpyrifos 4E may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above-listed foliar pests. For best results, use the listed rate of Quali-Pro Chlorpyrifos 4E per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See **SPRINKLER IRRIGATION** section for further information. For effective control of spider mites when large numbers of eggs are present, apply a second spray 3 to 5 days after initial treatment to control newly hatched nymphs.

For best results on bollworms and budworms, it is suggested that fields be scouted twice per week and treatments made when worms are 1/4 inch or less in length.

**RESTRICTIONS:** Do not apply within 14 days before harvest or make more than 3 applications per season. Do not apply more than 6 pints (3

*lbs. a.i. chlorpyrifos per acre per season. Do not make a second application within 10 days of the first application. Do not allow meat or dairy animals to graze in treated areas. Do not feed gin trash or treated forage to meat or dairy animals. The maximum single application rate is 1 lb. a.i. chlorpyrifos per acre.*

## CRANBERRIES

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E by application as a broadcast foliar spray to control brown spanworm, cranberry fruitworm, cranberry weevil, cutworms, fireworms, and *Sparganothis* fruitworm at the rate of 3 pints (1.5 lbs. a.i.) per acre. Mix the specified dosage in enough water to ensure thorough coverage and apply no less than 5 gallons of spray per acre when using aerial equipment or no less than 15 gallons of spray per acre when using ground equipment. For weevil control, apply once at flower bud development (late May, early June) and, if weevils are present, once after 100% bloom (early to mid-July). For other insects, treat when field counts indicate damaging insect populations are developing or present. Apply only after the winter flood has been removed. To avoid pesticide contamination of flood water, make no applications while bogs are flooded.

Quali-Pro Chlorpyrifos 4E may also be applied through sprinkler irrigation systems to control the above-listed pests. For best results, use the listed rate of Quali-Pro Chlorpyrifos 4E per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See **SPRINKLER IRRIGATION** section for further information.

**RESTRICTIONS:** Do not make more than two applications per year or apply within 60 days before harvest. Do not make a second application within 10 days of the first application. The maximum single application rate is 1.5 lbs. a.i. chlorpyrifos per acre.

## FIGS

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 days unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E at the rate of 2 quarts (2 lbs. a.i.) per acre for control of dried fruit beetle by application in sufficient water to the soil surface, followed by incorporation into the top 3 inches of soil. Apply to fig orchard soil as a dormant application in late winter prior to bee-tee emergence and prior to leaf formation. Use a spray volume of 10 gallons per acre or more and apply as a broadcast spray to the soil surface using ground equipment. On the day of treatment, incorporate into the top 3 inches of soil using suitable equipment.

**RESTRICTIONS:** Make only one application per year. Do not apply within 7 months (217 days) of harvest. Based on available residue data, use of Quali-Pro Chlorpyrifos 4E on figs is restricted to California. The maximum single application rate is 2 lbs. a.i. chlorpyrifos per acre.

## GRAPES

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E as a soil-surface application for control of grape root borer by application just before the pest emerges from the soil. Mix 4 1/2 pints (2.25 lbs. a.i.) of Quali-Pro Chlorpyrifos 4E with 100 gallons of water and apply 2 quarts of the diluted spray mixture to the soil surface on a 15-square-foot area around the base of each vine. Do not allow spray to contact fruit or foliage. Quali-Pro Chlorpyrifos 4E may also be used as a prebloom application to control climbing cutworm. Apply one quart per acre as a spray drench ground application using a minimum spray volume of 25 gallons per acre. Do not use in conjunction with soil-surface application for grape borer control.

**RESTRICTIONS:** Do not make more than one application per season or apply within 35 days before harvest. Based upon available residue data, the use of Quali-Pro Chlorpyrifos 4E in grapes is restricted to states east of the Rocky Mountains/Continental Divide. Do not make more than one application per season of Quali-Pro Chlorpyrifos 4E or other product containing chlorpyrifos. The maximum single application rate for soil-surface application is 2.12 lbs. a.i. chlorpyrifos per 100 gallons. The maximum single application rate for prebloom application is 1 lb. a.i.

### LEGUME VEGETABLES (SUCCULENT OR DRIED) (EXCEPT SOYBEAN)

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

CROP	SPECIFIC DIRECTIONS
Legume Vegetables including adzuki bean, asparagus bean, bean, blackeyed pea, broad bean (dry and succulent), catjang, chickpea, Chinese longbean, cowpea, crowder pea, dwarf pea, edible pod pea, English pea, fava bean, field bean, field pea, garbanzo bean, garden pea, grain lupin, green pea, guar, hyacinth bean, jackbean, kidney bean, lablab bean, lentil, lima bean, moth bean, mung bean, navy bean, pea, pigeon pea, pinto bean, rice bean, runner bean, snap bean, snow pea, southern pea, sugar snap pea, sweet lupin, sword bean, tepary bean, urd bean, wax bean, white lupin, white sweet lupin, yardlong bean	<p>For control of seed maggots, apply 2 pints/acre as a preplant broadcast application. Apply Quali-Pro Chlorpyrifos 4E in a minimum of 10 gpa of spray to the soil surface using suitable ground equipment. Quali-Pro Chlorpyrifos 4E must be incorporated in the top 1 to 3 inches of soil to improve the activity against seed maggots.</p> <p>To apply an at-plant T-band application, apply 1.8 fl. oz. of Quali-Pro Chlorpyrifos 4E per 1000 feet of row at 30-inch row spacing. To achieve shallow incorporation, apply the spray in a 3- to 5-inch wide band over the row behind the planter shoe and in front of the press wheel to achieve shallow incorporation. Mix the specified dosage in the table below in a minimum of 10 gpa of spray and apply to the soil surface as a ground spray. Equivalent rates of insecticide spray required per 100 feet of row for listed row spacing are given in the table below. Incorporate Quali-Pro Chlorpyrifos 4E into the top ½ to 1 inch of soil to improve activity against seed maggots.</p>

Spray Volume Per Acre (Gallons)	fl. oz. of Spray Volume per 100 Feet of Row			
	30-inch	28-inch	24-inch	22-inch
10	7.3	6.9	5.9	5.4
15	11	10.3	8.8	8.1
20	14.7	13.7	11.8	10.8

**RESTRICTIONS:** Do not make more than one application of Quali-Pro Chlorpyrifos 4E per year. Do not apply Quali-Pro Chlorpyrifos 4E at-plant if the field was treated with a preplant-incorporated treatment of Quali-Pro Chlorpyrifos 4E. Do not apply more than 2 pints of Quali-Pro Chlorpyrifos 4E (1 lb. a.i. chlorpyrifos) per acre. Not for use in Mississippi.

### MINT (PEPPERMINT AND SPEARMINT)

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E by application as a broadcast foliar spray to control cutworms at the rate of 2 to 4 pints (1 to 2 lbs. a.i.) per acre and mint root borer at the rate of 4 pints per acre. Mix the specified dosage in water to give no less than 10 gallons of spray per acre and apply using ground spray equipment. For cutworm control, treat during May and June when field counts indicate damaging insect populations are developing or present. When larvae are less than 3/4 inch in length, use the 2-pint rate. When larvae are 3/4 inch or more in length, use the higher rate. For mint root borer control, apply postharvest when field counts indicate damaging insect populations are developing or present.

Follow treatment with approximately 1 acre-inch of sprinkler irrigation immediately after application to incorporate the insecticide into the soil.

Quali-Pro Chlorpyrifos 4E may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the



above-listed pests. For best results, use the listed rate of Quali-Pro Chlorpyrifos 4E per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See **SPRINKLER IRRIGATION** for further information.

**RESTRICTIONS:** Do not apply within 90 days before harvest. Make only one application during the growing season. Do not make more than one preplant-incorporated application in the spring. Do not use in conjunction with a broadcast foliar application for cutworm control. Make only one postharvest application per season. The maximum single application rate is 2 lbs. a.i. chlorpyrifos per acre. Not for use in Mississippi.

### NECTARINES, PEACHES, ALMONDS (TRUNK SPRAY OR PREPLANT DIP)

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 days unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E for the control of peach tree borer in almonds, nectarines, and peaches by application as a trunk spray before newly hatched borers enter the trees. Mix 3 quarts (3 lbs. a.i.) of Quali-Pro Chlorpyrifos 4E with 100 gallons of water and apply as a coarse, low-pressure spray to give uniform coverage of tree trunks. Thoroughly wet all bark areas from ground level to scaffold limbs. Do not allow spray to contact fruit. Consult your State Agricultural Experiment Station or Extension Service specialist's written recommendations for proper time to treat in your area.

Quali-Pro Chlorpyrifos 4E may also be used as a preplant dip application for non-bearing peach and nectarine trees (only) at the equivalent application rate of 3 quarts (3 lbs. a.i.) per 100 gallons of water for control of peach tree borer. Dip trees several inches above the grafting bud scar and plant immediately or allow to dry before returning to storage. Do not allow peach trees to remain in contact with the dip solution.

**RESTRICTIONS:** Make only one application per season. Do not apply within 14 days before harvest. Do not allow meat or dairy animals to graze in treated orchards. Not for use in Mississippi.

### ONIONS (DRY BULB)

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

**At-Plant Soil Drench Application:** Use Quali-Pro Chlorpyrifos 4E to control onion maggot by application as an in-furrow drench. Apply Quali-Pro Chlorpyrifos 4E at the rate of 1.1 fluid ounces per 1000 linear feet of row at an 18-inch row spacing. Use a minimum of 40 gallons of total drench per acre. Incorporate to a depth of 1 to 2 inches. Do not make more than one application per year following these instructions.

**Post-Plant Soil Drench Application:** Apply one quart (1 lb. a.i.) per acre as an early-season directed spray to the base of onion seedlings or transplants during peak onion maggot egg laying. Use a minimum of 100 gallons per acre for thorough wetting.

**RESTRICTIONS:** Do not make more than two applications per year (at-plant plus post-plant). Do not harvest within 60 days of application. The maximum single application rate is 0.03 lb. a.i. chlorpyrifos per 1000 feet of row. Do not aerially apply this product in Mississippi.

### PEANUTS

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

For suppression of wireworms, apply Quali-Pro Chlorpyrifos 4E at a rate of 4 pints (2 lbs. a.i.) per acre as a preplant broadcast spray to the soil surface, followed by immediate soil incorporation to a depth of 3 to 4 inches. Use a minimum of 10 gallons of total spray per acre.

**RESTRICTIONS:** Do not make more than one application per season. The maximum single application rate is 2 lbs. a.i. chlorpyrifos per acre. Do not harvest within 21 days after treatment. Do not feed treated peanut forage or hay to meat or dairy animals. Aerial application to peanuts is prohibited in Mississippi.

**PEARS  
(California, Oregon, and Washington)**

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Apply 4 pints (2 lbs. a.i.) per acre in 100 to 400 gallons of spray per acre to control codling moth. Apply using an airblast speed sprayer or other suitable ground equipment. For this use, do not make more than one postharvest application prior to dormancy per year, do not harvest or use treated fruit for food or feed, and do not allow meat or dairy animals to graze in treated orchards. If unauthorized entry into a treated orchard cannot be prevented, then the orchard shall be posted with appropriate signs according to the Worker Protection Standard while treated, unharvested fruit remains on the tree.

**SORGHUM – GRAIN SORGHUM (MILO)**

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E for control of the following pests at the dosage indicated:

<b>PEST</b>	<b>QUALI-PRO CHLORPYRIFOS 4E</b>	<b>SPECIFIC DIRECTIONS</b>
Sorghum midge	1/2 pint/acre (0.25 lb. a.i./acre)	Apply when 30% to 50% of the seed heads are in bloom. Repeat at 3-day intervals if necessary.
Grasshoppers, Yellow sugar cane aphid and other aphids	1/2 – 1 pint/acre (0.25 – 0.5 lb. a.i./acre)	
Greenbug	1/2 – 2 pints/acre (0.25 – 1 lb. a.i./acre)	For infestations of greenbug that are difficult to control, use a higher dosage within the indicated rate range.
Chinch bugs, Lesser cornstalk borer	1 – 2 pints/acre (0.5 – 1 lb. a.i./acre)	Apply as a directed spray toward the base of the plant using power-operated ground spray equipment with sufficient water to ensure coverage of an 8- to 12-inch band centered on the row. On plants less than 6 inches high, apply an 8- to 12-inch band over the row. Do not reduce the dosage for banded or directed applications. Concentrate the full labeled dosage rate in the treated zone.
Webworms	1 pint/acre (0.5 lb. a.i./acre)	
Armyworms, Corn earworm, Cutworms	1 – 2 pints/acre (0.5 – 1 lb. a.i./acre)	
European and Southwestern corn borer	1 1/2 – 2 pints/acre (0.75 – 1 lb. a.i./acre)	

Mix the specified dosage in enough water to ensure thorough coverage and apply using suitable aerial or ground spray equipment. To minimize chemical injury, do not apply Quali-Pro Chlorpyrifos 4E to drought-stressed grain sorghum within 3 days following irrigation or rain except where the product is applied in irrigation water.

Quali-Pro Chlorpyrifos 4E may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the

above-listed foliar pests. For best results, use the listed rate of Quali-Pro Chlorpyrifos 4E per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See **SPRINKLER IRRIGATION** section for further information.

**RESTRICTIONS:** *The treated crop is not to be used for grain, forage, fodder, hay, or silage within 30 days after application of 1 pint of Quali-Pro Chlorpyrifos 4E per acre or within 60 days after application of rates above 1 pint (0.5 lb. a.i.) per acre. Do not treat sweet varieties of sorghum. Do not apply more than 3 pints (1.5 lbs. a.i.) of Quali-Pro Chlorpyrifos 4E per acre per season. Do not make more than 3 applications per season. Do not make a second application within 10 days of the first application. Do not apply by air in Mississippi.*

## SOYBEANS

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

For use to control armyworms, bean leaf beetle, corn earworm, cutworms, European corn borer, grasshoppers, green cloverworm, lesser cornstalk borer, Mexican bean beetle, saltmarsh caterpillar and other woollybears, southern green stink bug, spider mites, and velvetbean caterpillar.

### Soil Treatment

Use Quali-Pro Chlorpyrifos 4E at the rate of 1 to 2 pints (0.5 to 1 lb. a.i.) per acre to control cutworms and lesser cornstalk borer. Mix the specified dosage in a minimum of 10 gallons of spray per acre and apply to the soil surface using suitable ground spray equipment. Equivalent rates of insecticide spray required per 100 feet of row for various row spacing are given in the accompanying table. For at-plant treatments, apply the insecticide over the row in a 4- to 6-inch band in front of the planter shoe or press wheel or after the press wheel, followed by a drag chain for light incorporation. Do not apply as an in-furrow treatment. For postemergence rescue treatments, apply as a directed spray in a 9- to 12-inch band at the base of the plant. To plants under 6 inches high, apply over-the-top in a 6- to 12-inch band. Treat when field counts or conditions indicate that pests are or may become a problem.

**Fluid Ounces of Spray Required Per 100 Feet of Row for Various Row Spacing**

VOLUME OF SPRAY PER ACRE	36"	32"	28"	24"
10 gallons	8.8	7.9	6.9	5.9
15 gallons	13.2	11.8	10.3	8.8
20 gallons	17.6	15.7	13.7	11.8

### Foliar Treatment

Use Quali-Pro Chlorpyrifos 4E at the following rate by application in sufficient water to ensure thorough coverage of treated plants:

PEST	QUALI-PRO CHLORPYRIFOS 4E
European corn borer, Southern green stink bug	2 pints/acre (1 lb. a.i./acre)
Bean leaf beetle, Cutworms, Corn earworm, Saltmarsh caterpillar and other woollybears, Soybean aphid	1 – 2 pints/acre (0.5 – 1 lb. a.i./acre)
Mexican bean beetle, Armyworms	1 – 1 1/2 pints/acre (0.5 – 0.75 lb. a.i./acre)
Velvetbean caterpillar, Grasshoppers, Green cloverworm, Spider mites	1/2 – 1 pint/acre (0.25 – 0.5 lb. a.i./acre)

Apply as a broadcast spray using either aerial or ground equipment when field counts indicate damaging insect populations are developing or present; re-treat as necessary to maintain control. Mix the specified dosage in a minimum of 15 gallons of spray per acre for ground spray

equipment or 5 gallons of spray per acre for aircraft equipment. For effective control of spider mites when large numbers of eggs are present, apply a second spray 3 to 5 days after initial treatment to control newly hatched nymphs. On determinate soybeans, do not apply more than one application after pod set.

Quali-Pro Chlorpyrifos 4E insecticide may also be applied through sprinkler irrigation systems as a postemergence broadcast application to control the above-listed foliar pests. For best results, use the listed rate of Quali-Pro Chlorpyrifos 4E per acre. Maintain vigorous tank agitation to assure uniformity of the application throughout the injection period. See **SPRINKLER IRRIGATION** section for further information.

**RESTRICTIONS:** Do not apply more than 6 pints (3 lbs. a.i.) of Quali-Pro Chlorpyrifos 4E per acre per season. Do not make more than 3 applications per season. Do not apply last treatment within 28 days before harvest nor apply last two treatments closer than 10 days apart. Do not allow livestock to graze in treated areas or otherwise feed treated soybean forage, hay, and straw to meat or dairy animals. Not for use in Mississippi.

## STRAWBERRIES

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

**Preplant Incorporation Treatment:** Use Quali-Pro Chlorpyrifos 4E insecticide at the following rates by application in sufficient water to the soil surface and incorporate into the soil in the spring for protection of strawberries during the following year:

PEST	QUALI-PRO CHLORPYRIFOS 4E
Garden symphylans, Grub	4 pints/acre (2 lbs. a.i./acre)

**Foliar Application:** Use Quali-Pro Chlorpyrifos 4E by application as a broadcast foliar spray to control strawberry bud weevil (clipper) at the rate of 1 quart per acre. Apply in a minimum of 40 gallons of spray per acre when buds first appear and 10 to 14 days later. Do not apply after berries start to form or when berries are present. Do not tank mix Quali-Pro Chlorpyrifos 4E with pesticides, surfactants, or fertilizer formulations unless prior use has shown the combination non-injurious under your current conditions of use. Phytotoxicity may occur when Quali-Pro Chlorpyrifos 4E is applied to strawberries experiencing high temperature and drought stress.

**Postharvest Application:** For strawberry crown moth, apply 1 quart (1 lb. a.i.) per acre as a directed spray to crown of strawberry plants immediately after harvest and after plants are topped. Use a minimum spray volume of 100 gallons per acre and repeat application if required 14 to 18 days later. Do not sprinkle irrigate for one week following application.

**RESTRICTIONS:** For prebloom use only, do not apply after berries start to form or when berries are present. Do not apply within 21 days before harvest. For preplant application, do not make more than one application per year. For foliar and postharvest applications, do not make more than two applications per year. Do not make a second application within 10 days of the first foliar application and within 14 days for postharvest application. The maximum single application rate is 2 lbs. a.i. chlorpyrifos per acre and 1 lb. a.i. chlorpyrifos per acre for foliar and postharvest application. Not for use in Mississippi.

## SUGAR BEETS

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

**Soil Treatment (At-Planting or Preplant Incorporated):** To reduce feeding damage from early-season insects such as cutworms, apply at planting or as a preplant treatment and incorporate to a depth of 1 to 2 inches. Do not apply as an in-furrow treatment. Apply 1 pint of Quali-Pro Chlorpyrifos 4E per planted acre to a 10-inch wide band centered over the row for furrows 30 inches apart. (For rows 30 inches apart, this is equivalent to 9.2 fl. oz. of Quali-Pro Chlorpyrifos 4E per 10,000 feet of row.) For other row widths, adjust the spray volume per planted acre in proportion to the length of row actually treated.

**Postemergence Treatment:** Apply specified rate as a broadcast or banded foliar spray. Treat when field counts indicate that damaging insect populations are developing or present.

**Broadcast Application:** Apply the specified dosage in water using 2 to 5 gallons per acre of finished spray when using aerial spray equipment or 10 to 30 gallons per acre when using ground spray equipment.

**Banded Foliar Spray:** Apply the specified rate within the band using a minimum of 7 gallons of spray volume in a 5- to 7-inch wide band centered over the row. Do not reduce the rate for band applications. Concentrate the full labeled dosage rate (see band rates in table below) in the treated zone. For best results, band-applied treatments should be lightly incorporated, either mechanically or with irrigation.

TARGET PESTS	QUALI-PRO CHLORPYRIFOS 4E	
	BROADCAST (PINT/ACRE)	BAND (PINT/ACRE)
Grasshoppers	1/2 – 1 (0.25 – 0.5 lb. a.i.)	–
Spider mites	1 (0.5 lb. a.i.)	2/3 (0.33 lb. a.i.)
Tarnished plant (Lygus)	1 (0.5 lb. a.i.)	–
Fall armyworms, Yellowstriped armyworm, Webworms	1 – 2 (0.5 – 1 lb. a.i.)	2/3 – 1 1/3 (0.33 – 0.66 lb. a.i.)
Beet armyworm	1 1/2 – 2 (0.75 – 1 lb. a.i.)	1 – 1 1/3 (0.5 – 0.66 lb. a.i.)
Cutworms, Flea beetle adults	2 (1 lb. a.i.)	1 1/3 (0.66 lb. a.i.)
Sugarbeet root maggot adults	1/2 – 1 (0.25 – 0.5 lb. a.i.)	–
Sugarbeet root maggot larvae	–	1 1/3 – 2 (0.66 – 1 lb. a.i.)
Sugarbeet root maggot larvae	2 (1 lb. a.i.)	1 1/3 – 2 (0.66 – 1 lb. a.i.)
Aphids*	1-2 (0.5-1 lb. a.i.)	–

\*Not for use in California on aphids.

For grasshopper control, the low rate will control small nymphs (1st through 3rd instar). For sugarbeet root maggot adults, apply anytime from 7 days before until 3 days after peak adult emergence in order to target adults present at time of application based on local field trap monitoring. For sugarbeet root maggot larvae, use as primary treatment to control root maggot larvae. Base application timing on local field trap monitoring. Apply anytime from 7 days before until 3 days after peak adult emergence. Use as a supplemental postemergence treatment following an at-plant insecticide application for control of root maggot larvae. Base application timing on local field trap monitoring. Apply anytime from 7 days before until 3 days after peak adult emergence. To prevent potential development of insecticide resistance in sugarbeet root maggot, producers are encouraged to take the following steps: (1) avoid making more than 2 applications of Quali-Pro Chlorpyrifos 4E per season when adults are active; (2) If an organophosphate insecticide was applied at planting, make no more than one postemergence application of Quali-Pro Chlorpyrifos 4E when adults are active.

**RESTRICTIONS:** Do not apply within 30 days before harvest of beet roots and tops. Do not make more than 3 applications per season. Do not apply more than a total of 6 pints (3 lbs. a.i.) per acre per season. The maximum single application rate is 1 lb. a.i. chlorpyrifos per acre. Do not make a second application within 10 days of the first application. Do not allow livestock to graze in treated areas nor harvest treated beet tops as feed for meat or dairy animals within 30 days after last treatment. Not for use in Mississippi.

#### SUNFLOWERS

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

For use to control cutworms, sunflower beetle larvae and adults, stem weevil, sunflower moth, banded sunflower moth, woollybears, seed weev-

wil, and grasshoppers.

### Preplant Incorporation Treatment

Use Quali-Pro Chlorpyrifos 4E insecticide at the following rates by application in sufficient water to the soil surface and incorporate into the soil:

PEST	QUALI-PRO CHLORPYRIFOS 4E
Cutworms	2 – 4 pints/acre (1 – 2 lbs. a.i./acre)

Use listed rate in not less than 10 gallons of water per acre and apply as a broadcast spray to the soil surface using suitable power-operated ground spray equipment. On the same day of treatment, incorporate the insecticide into the top 2 to 4 inches of soil using a disc, field cultivator, or equivalent equipment.

### Postemergence Treatment

Use Quali-Pro Chlorpyrifos 4E for control of the following pests at the dosage indicated by application in sufficient water to ensure thorough coverage of treated plants:

PEST	QUALI-PRO CHLORPYRIFOS 4E
Cutworms	2 – 3 pints/acre (1 – 1.5 lbs. a.i./acre)
Sunflower beetle larvae and adults, Stem weevil, Sunflower moth, Banded sunflower moth, Woollybears, Seed weevil	1 – 1 1/2 pints/acre (0.5 – 0.75 lb. a.i./acre)
Grasshoppers	1 pint/acre (0.5 lb. a.i./acre)

Apply as a broadcast spray using either aerial (fixed-wing or helicopter) or power-operated ground spray equipment when field counts indicate that pests are or may become a problem. For cutworm control, a second treatment may be made 7 to 10 days later if needed. For stem weevil control, optimal treatment time is within 5 to 7 days after adult weevils begin to appear. For sunflower moth control, make first application during early 1- to 5-percent bloom stage. A second treatment may be made 10 days later if needed. For seed weevil control, treat when field counts indicate there are 10 to 12 adults per plant for oil crops and 1 to 3 adults per plant on confectionery crops.

Additional treatments should be made at successive 10-day intervals if field counts indicate need to re-treat. For sunflower beetle larvae or adult control, treat when field counts indicate there are 10 larvae or 1 to 2 adults per seedling. Additional treatments may be made at successive 10-day intervals if field counts indicate need to re-treat.

**RESTRICTIONS:** Do not apply more than 6 pints (3 lbs. a.i.) of Quali-Pro Chlorpyrifos 4E per acre per season. Do not make more than 3 applications per season. Do not apply within 42 days before harvest. Do not make a second application within 10 days of the first application. Do not allow livestock to graze in treated areas. The maximum single application rate is 2 lbs. a.i. chlorpyrifos per acre for preplant incorporation and 1.5 lbs. a.i. chlorpyrifos per acre for postemergence broadcast treatment. Not for use in Mississippi.

### SWEET POTATOES

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E to reduce the feeding damage caused by populations of *Conderus* wireworm, *Systema* flea beetle, and the sweet potato flea beetle. Apply at the rate of 4 pints (2 lbs. a.i.) per acre as a broadcast (overall) spray to the soil surface followed by incorporation. Mix the specified dosage with enough water to obtain uniform coverage and apply as a coarse spray using suitable ground spray equipment. Incorporate the insecticide to a depth of 4 to 6 inches as soon as possible after application by using a rotary hoe, disc cultivator, or other suit-

able incorporation equipment. Plant the crop in the usual manner no later than 14 days after treatment (any delay in planting will reduce the length of time that Quali-Pro Chlorpyrifos 4E will protect against feeding damage). Quali-Pro Chlorpyrifos 4E will not control false wireworms or whitefringed beetle or other grubs that attack sweet potatoes.

**RESTRICTIONS:** Do not make more than one application per season. Do not harvest within 125 days of treatment. The maximum single application rate is 2 lbs. a.i. chlorpyrifos per acre. Do not aerially apply this product in Mississippi.

## TOBACCO

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E for preplant treatment to control larvae of cutworms, flea beetles, mole crickets, root maggots, and wireworms. Apply 2 pints (1 lb. a.i.) of Quali-Pro Chlorpyrifos 4E per acre in not less than 10 gallons of water as a broadcast (overall) spray to the soil surface one week before transplanting. Immediately following application, incorporate the insecticide into the soil to a depth of 2 to 4 inches using suitable equipment.

To control the above insects and low to moderate populations of rootknot nematodes in North Carolina, South Carolina, and Virginia, use Quali-Pro Chlorpyrifos 4E at the rate of 2 pints (1 lb. a.i.) per acre. To control the above insects and moderate populations of rootknot nematodes in all tobacco-growing regions, use Quali-Pro Chlorpyrifos 4E in a tank mix with Nemacur® 3 at the rate of 2 quarts (2 lbs. a.i.) of Quali-Pro Chlorpyrifos 4E plus 4 quarts of Nemacur 3 nematicide per acre. Read and carefully follow all applicable directions, restrictions, and precautions on labeling for Nemacur 3 used in combination with Quali-Pro Chlorpyrifos 4E.

Apply the specified dosage in not less than 10 gallons of water as a broadcast (overall) spray to the soil surface 24 to 48 hours before bedding and transplanting. Immediately following application, incorporate into the soil surface 24 to 48 hours before bedding and transplanting. Immediately following application, incorporate into the soil to a depth of at least 4 inches using suitable equipment. Where the nematode species *Meloidogyne arenaria* or *M. javanica* are present or high populations of *M. incognita*, apply Telone® II soil fumigant at the listed label rate.

**RESTRICTIONS:** Do not apply more than 2 pints (1 lb. a.i.) per application. Do not make more than one application per season. Do not apply more than 2 pints (1 lb. a.i.) per acre per season. Do not aerially apply this product in Mississippi.

## TREE FRUITS (DORMANT/DELAYED-DORMANT SPRAYS)

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 days unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E as a dormant or delayed-dormant spray at the rates indicated to control the following insects on the crops listed. While Quali-Pro Chlorpyrifos 4E may be used without oil, oil is recommended to control additional pests such as European red mite and brown almond mite. Applications may be made on pears after harvest. See specific use directions below.

CROP	PEST	QUALI-PRO CHLORPYRIFOS 4E PER 100 GALLONS OF SPRAY*
Apples (1)	Rosy apple aphid, San Jose scale, Lygus, <i>Pandemis</i> leafroller, Climbing cutworms, Obliquebanded leafroller	1/2 – 1 pint (0.25 – 0.5 lb. a.i.)  (Use a minimum of 1.5 pints/acre)
Pears	San Jose scale, Climbing cutworms, Pear psylla adults	
Plums, Prunes	San Jose scale, Mealy plum aphid, Climbing cutworms, Peach twig borer	
Almonds, Peaches, Nectarines	San Jose scale, Peach twig borer, Climbing cutworms	

\* Based on 200 to 600 gallons per acre as a dilute spray.

(1) Only one application of any chlorpyrifos-containing product may be made per year. The application can either be a prebloom dormant/dormant-delayed spray to the canopy or the trunk, OR a postbloom application to the lower 4 feet of the trunk. For postbloom application instructions and restrictions for apple, see the **APPLE (TREE TRUNK)** section.

For dilute spray, tank mix the specified dosage with 1 to 2 gallons of a petroleum spray oil recommended for dormant use in 100 gallons of water and spray the entire tree by application to runoff using suitable ground spray equipment. (See **Additional Precautions Specific to California** section for use in California.)

For low-volume (concentrate) sprays (less than 200 gallons of spray mixture per acre), use the same amount of Quali-Pro Chlorpyrifos 4E as for a dilute spray and apply in a manner that will ensure thorough coverage of the trees. Use the higher dosage of Quali-Pro Chlorpyrifos 4E for severe infestations. Use oil as recommended by your State Agricultural Experiment Station or Extension Service specialist.

**PRECAUTIONS:** Because cold or dry conditions may cause Quali-Pro Chlorpyrifos 4E plus oil sprays to infuse trees resulting in bud damage or drop, do not apply until winter rains or irrigation has replenished soil moisture such that bark and twigs are not desiccated. Do not use more than 4 pints of Quali-Pro Chlorpyrifos 4E per acre.

**Additional Precautions Specific to California:** Use a minimum of 100 gallons of total spray volume per acre. Do not use more than 4 gallons of spray oil per acre on almonds, and 6 gallons per acre on peaches and nectarines. Do not use any adjuvants or surfactants in addition to or as a substitute for a petroleum spray oil in a tank mix with Quali-Pro Chlorpyrifos 4E. Do not apply on almonds in the following counties in California: Butte, Colusa, Glenn, Solano, Sutter, Tehama, Yolo, and Yuba. Do not use more than 1% dormant oil in almond orchards less than 4 years old.

**RESTRICTIONS:** *Make only one application during the dormant/delayed-dormant season, applying no more than 4 pints (2 lbs. a.i.) per acre per season. Do not allow meat or dairy animals to graze in treated orchards. Only one application of any chlorpyrifos-containing product may be made per year. The application can either be a prebloom dormant/dormant-delayed spray to the canopy or the trunk, OR a postbloom application to the lower 4 feet of the trunk. For postbloom application instructions and restrictions for apple, see the **APPLE (TREE TRUNK)** section.*

*Do not make a soil or foliar application within 10 days of a dormant/delayed-dormant application of chlorpyrifos to the orchard.*

## TREE NUTS

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

**Foliar Spray:** Use Quali-Pro Chlorpyrifos 4E at the dosage indicated by application as a foliar spray to control pests listed in the following table. Mix the required dosage in sufficient water to ensure thorough and complete coverage of the foliage and crop, and apply as a concentrate or dilute spray using conventional, power-operated spray equipment. For dilute sprays applied to tree nut crops, mix the required dosage in sufficient water to allow for spray to runoff. For concentrate sprays, apply an equivalent amount of Quali-Pro Chlorpyrifos 4E per acre. Treat when pests appear or in accordance with local conditions. Insect control by aerial application may be less than control by ground application because of less coverage. Consult your State Agricultural Experiment Station, certified pest control advisor, or Extension Service specialist for specific use information in your area. To avoid contamination of irrigation floodwaters, do not flood irrigate within 24 hours following an application of Quali-Pro Chlorpyrifos 4E. Quali-Pro Chlorpyrifos 4E is highly toxic to bees exposed to direct treatment. Do not apply when bees are actively foraging in the treated area.

## ALMONDS, FILBERTS, PECANS, WALNUTS

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E at the rates indicated to control the listed pests.



CROP	PEST	DOSAGE
Almonds	Navel orangeworm, Peach twig borer, San Jose scale	4 pints/acre (2 lbs. a.i./acre)
Filberts	Eyespotted bud moth, Filbert aphid, Filbert leafroller, Filbert worm, Oblique-banded leafroller, Omnivorous leaflier, Winter moth	3 – 4 pints/acre (1.5 – 2 lbs. a.i./acre)
Walnuts	Codling moth, Walnut scale, Walnut husk fly	4 pints/acre (2 lbs. a.i./acre)
Pecans	Pecan nut casebearer, Fall webworm	1 1/2 – 4 pints/acre (0.75 – 2 lbs. a.i./acre)
	<i>Phylloxera</i> spp. <sup>1</sup> , Black pecan aphid, Hickory shuckworm <sup>2</sup> , Pecan leaf scorch mite (suppression) <sup>3</sup> , Ant species (except fire ants, carpenter ants, harvester ants, and pharaoh ants) <sup>4</sup>	2 – 4 pints/acre (1 – 2 lbs. a.i./acre)
	Yellow pecan aphid <sup>5</sup> , Black margined aphid <sup>5</sup> , Spittlebugs	1 – 4 pints/acre (0.5 – 2 lbs. a.i./acre)

<sup>1</sup> For best *Phylloxera* spp. control, make 2 applications at 10-day intervals using a minimum of 1 pint of Quali-Pro Chlorpyrifos 4E per acre starting at bud swell.

<sup>2</sup> For best results, make 2 applications 10-14 days apart.

<sup>3</sup> To suppress pecan leaf scorch mite, use a preventative program.

<sup>4</sup> For ant control, apply as an orchard floor spray. Do not apply where weed growth or other obstructions prevent uniform coverage of the orchard floor.

<sup>5</sup> For control of yellow pecan aphid and black margined aphid, apply in tank mix combination with the listed rate of a pyrethroid insecticide labeled for control or suppression of these aphids.

**Dormant or Delayed-Dormant Spray:** For control of additional pests such as European red mite, it is recommended to mix Quali-Pro Chlorpyrifos 4E with oil, although it can be used without oil. Using conventional, power-operated spray equipment, apply as a concentrate or dilute spray. For dilute sprays (greater than 200 gpa), use sufficient spray volume to saturate tree foliage, but not to point of runoff. For concentrate sprays (less than 200 gpa), evenly distribute an equivalent amount of product per acre.

**Precautions:** Cold or dry conditions may cause this product and oil mixtures to infuse into trees, resulting in bud damage or bud drop. Do not apply until winter rains or irrigation has replenished soil moisture so that bark and twigs are not desiccated. Do not flood irrigate within 24 hours of application of this product to avoid contamination of irrigation tail waters.

**RESTRICTIONS:** Make no more than 3 foliar applications per season on almonds, filberts, and pecans and no more than 2 applications per season on walnuts. Make no more than one application of chlorpyrifos during the dormant season. Do not make a soil or foliar application of Quali-Pro Chlorpyrifos 4E or products containing chlorpyrifos within 10 days of a dormant/delayed-dormant application. Do not apply within 14 days of harvest of almonds, filberts, and walnuts, or 28 days of harvest of pecans. Do not allow livestock to graze in treated orchards. Do not apply more than 8 pints (4 lbs. a.i.) per acre per season as a foliar spray. Do not make a second application within 10 days of the first application. Do not apply more than 4 pints of Quali-Pro Chlorpyrifos 4E (2 lbs. a.i. chlorpyrifos) per acre per season as a dormant/delayed-dormant application. Do not aerially apply this product in Mississippi.

**CALIFORNIA RESTRICTIONS:** Do not use more than 1% dormant oil and/or penetrating surfactants in almond orchards less than 4 years old. Use a minimum of 100 gpa of total spray volume. Use up to 2% supreme oil with no more than 4 gpa on almonds. In orchards with high overwintering populations of European red mite or brown almond mite, use the higher spray volumes that allow for the use of higher per acre rates of oil. Do not use any adjuvants or surfactants in addition to or as a substitute for petroleum spray oil in a tank mix with this product. Do not apply on almonds during dormant and delayed dormant in the following counties in California: Butte, Colusa, Glenn, Solano, Sutter, Tehama, Yolo, and Yuba.

## ALMOND, PECAN, WALNUT ORCHARD FLOORS

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E to control ants (except fire ants, carpenter ants, harvester ants, and pharaoh ants) and pavement ant by applying the specified dose with ground application equipment that will uniformly apply the spray to the orchard floor. Use when ant activity becomes evident within the orchard.

Since worker ants cease most of their foraging activity at temperatures above 90° F, best results will be achieved with applications made at temperatures below 90° F at the time of application. Dosage of Quali-Pro Chlorpyrifos 4E and spray volume may vary depending on the irrigation method employed in the orchard as follows:

### **Ant Control in Sprinkler- or Drip-irrigated Orchards**

Apply Quali-Pro Chlorpyrifos 4E as a broadcast spray to the entire orchard floor using ground spray equipment at 4 to 8 pints (2 – 4 lbs. a.i.) per acre in 25 or more gallons of water. Use the high rate for heavy infestations and the low rate for light infestations. In orchards where ant activity is concentrated around the irrigation emitters, apply the high rate to a 6- to 8-foot band along the drip-irrigation line and the low rate to the rest of the orchard.

### **Ant Control in Flood-irrigated Orchards**

Apply Quali-Pro Chlorpyrifos 4E at 4 to 8 pints (2 – 4 lbs. a.i.) per acre in 25 or more gallons of water to the entire orchard floor using ground spray equipment. Apply the high rate to heavily infested areas and the low rate to lightly infested areas. Where ant colonies are abundant only in the berm areas, apply Quali-Pro Chlorpyrifos 4E at 8 pints (4 lbs. a.i.) per treated acre in 50 or more gallons of water to a 6- to 10-foot band along the treeline (berm).

Do not apply where weed growth or other obstructions would impede uniform coverage of the orchard floor. Mow or chemically control weeds before the application of Quali-Pro Chlorpyrifos 4E. Foliar applications of Quali-Pro Chlorpyrifos 4E may be made in addition to the orchard floor treatment.

**RESTRICTIONS:** Do not make more than 2 applications per season. Do not apply more than 8 pints (4 lbs. a.i.) per acre per season to the orchard floor. If the 8-pint rate is used, then a second application is not permitted. Do not apply the last treatment within 14 days of harvest. Do not allow livestock to graze in treated orchards. Do not allow spray to contact fruit or foliage. Do not make a second application within 10 days of the first application.

## TURFGRASS GROWN FOR COMMERCIAL SOD

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E to control the pests listed in the following table by application at the listed dosages. Dilute Quali-Pro Chlorpyrifos 4E in water and apply using suitable application equipment. For best results, turf should be moist at time of treatment.

PESTS	AMOUNT OF QUALI-PRO CHLORPYRIFOS 4E	
	1000 SQ. FT.	ACRE
Ants (except fire ants, carpenter ants, harvester ants, and pharaoh ants), Armyworms (such as beet, fall, yellowstriped), Chinch bugs, Crickets, Cutworms, Earwigs, European crane fly larvae, Fiery skipper, Gnats, Grasshoppers, Greenbug aphids, Green June beetle grubs, Leafhoppers, Lucerne moth, Millipedes, Mites (such as clover, Bermudagrass stunt, winter grain), Pillbugs, Springtails, Sod webworms (lawn moths) (1), Sowbugs	3/4 fl. oz.	1 qt. (1 lb. a.i.)
Billbug adults (such as bluegrass, Denver, hunting) (2)	3/4 – 1 1/2 fl. oz.	1 – 2 qts. (1 – 2 lbs. a.i.)
Annual bluegrass weevil (Hyperodes) (3), Black turfgrass ateniaus adults (4), Mole crickets (5)	1 1/2 fl. oz.	2 qts. (2 lbs. a.i.)
White grubs (such as black turfgrass ateniaus, European chafer, Japanese beetle larvae, and Northern and Southern masked chafers) (6)	1 1/2 – 3 fl. oz.	2 – 4 qts. (2 – 4 lbs. a.i.)

Numbers in parentheses (-) refer to **SPECIFIC USE DIRECTIONS** below.

#### **SPECIFIC USE DIRECTIONS:**

1. For sod webworms, watering or mowing of the treated area should be delayed for 12 to 24 hours after treatment.
2. For billbugs, spray early in the season just prior to or coinciding with first appearance of adults as recommended by your local Agricultural Extension Service specialist.
3. To control annual bluegrass weevil, spray suspected problem areas in mid-April and again in mid-May, or as recommended by your local Agricultural Extension Service specialist.
4. For black turfgrass ateniaus adults, spray early in the season as recommended by your local Agricultural Extension Service specialist. A repeat application may be needed 1 to 2 weeks later.
5. To control mole crickets in turfgrass, apply Quali-Pro Chlorpyrifos 4E through high-pressure injection or other suitable subsurface placement application equipment. Depending on the application equipment used, follow the manufacturer's recommendation for calibration and the volume of spray per acre needed to provide control or as recommended by your local Agricultural Extension Service specialist. For best results, apply when young nymphs are active.
6. For white grubs, spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by your local Agricultural Extension Service specialist. For best results, soil should be moist prior to treatment. For best results, immediately after spraying, irrigate the treated area with 1/2 to 1 inch of water to wash the insecticide into the thatch and underlying soil.

#### **VEGETABLES (BRASSICA, COLE CROPS)**

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 3 days for cauliflower and 24 hours for all other vegetables unless PPE required for early entry is worn.

Use Quali-Pro Chlorpyrifos 4E at the dosages indicated to control the pests listed in the following table. To avoid phytotoxicity in vegetables except Brussels sprouts, do not mix with other pesticide products or treat plants that are under extreme heat and drought stress.

CROP	PEST	DOSAGE QUALI-PRO CHLOR- PYRIFOS 4E	USE DIRECTIONS	RESTRICTIONS
Cauliflower – soil application	Root maggot	1.6 – 2.4 fl. oz./ 1000 linear ft. of row	For direct-seeded crops, apply the specified dosage in a water-based spray as a 4-inch wide band over the row at planting time. Shallow incorporation is necessary. Placement behind the planter shoe and in front of the press wheel is recommended. For transplanted crops, apply Quali-Pro Chlorpyrifos 4E as a water-based spray directed to the base of the plants immediately after setting. Use a minimum of 40 gallons of total spray per acre. Do not add any additional adjuvants, surfactants, or spreader stickers. Do not apply as a foliage application.	Do not apply more than 2 pints of Quali-Pro Chlorpyrifos 4E to cauliflower planted in 40-inch rows. Use proportional amounts for other row spacings not to exceed 4 pints of Quali-Pro Chlorpyrifos 4E per acre. Do not apply more than 2.6 pints (1.3 lbs. a.i.) of Quali-Pro Chlorpyrifos 4E per acre to broccoli, Brussels sprouts, cabbage, Chinese cabbage, collards, kale, kohlrabi, and turnips planted in 40-inch rows. Do not apply more than 4.5 pints (2.25 lbs. a.i.) of Quali-Pro Chlorpyrifos 4E per acre to these crops in 20-inch rows (or two rows per bed). Use proportional amounts for other row spacings not to exceed 4.5 pints (2.25 lbs. a.i.) of Quali-Pro Chlorpyrifos 4E per acre. Do not make more than one application per season within 30 days before harvest on broccoli, Brussels sprouts, cabbage, Chinese cabbage, collards, kale, kohlrabi, and turnips. Do not make more than one soil application per crop.
Broccoli, Brussels sprouts, Cabbage, Chinese cabbage, Collards, Kale, Kohlrabi, Turnips – soil application		1.6 – 2.75 fl. oz./ 1000 linear ft. of row		
Broccoli, Cabbage – soil application	Root aphid	1.2 fl. oz./ 1000 ft. of row for single row plantings, and 2.4 fl. oz./ 1000 linear ft. of row for double row plantings	Apply Quali-Pro Chlorpyrifos 4E in a water emulsion or with liquid fertilizer injected as a sidedress on each side of the row after plants are established. Avoid mechanical damage to crop roots. Use a minimum of 15 gallons of total spray volume per acre. See <b>MIXING DIRECTIONS</b> section of the label for mixing instructions with liquid fertilizers.	Do not make a foliar application within 10 days of a soil application.
Brussels sprouts – foliar application	Armyworms, Cabbage aphid, Cutworms, Imported cabbage worm, Striped flea beetle (adult)	1 – 2 pints/acre (0.5 – 1 lb. a.i./acre)	Apply Quali-Pro Chlorpyrifos 4E with conventional power-operated equipment in 20 to 150 gallons of water per acre. Apply when insects appear on foliage and at 10- to 14-day intervals or thereafter as needed. Consult your State Agricultural Station, Extension Service specialist, or integrated pest control advisor for proper time to treat in your area.	Do not make more than 3 applications per season. Do not apply within 21 days before harvest. Do not make a second application within 10 days of the first application. Do not make a foliar application within 10 days of a soil application.

**VEGETABLES (BRASSICA, COLE CROPS) (continued)**

<b>CROP</b>	<b>PEST</b>	<b>DOSAGE QUALI-PRO CHLOR- PYRIFOS 4E</b>	<b>USE DIRECTIONS</b>	<b>RESTRICTIONS</b>
Radishes – soil application	Root maggot	1.0 fl. oz./ 1000 linear ft. of row	Apply the specified dosage as a water-based drench in the seed furrows with the seed at planting time. Use a minimum of 40 gallons of total drench per acre.	Do not apply more than 5.5 pints (2.75 lbs. a.i.) of Quali-Pro Chlorpyrifos 4E per acre or make more than one soil application per season. Do not make a foliar application within 10 days of a soil application. Do not apply within 30 days of harvest.
Rutabagas – soil application	Root maggot	1.6 – 3.2 fl. oz./ 1000 linear ft. of row	Apply the specified dosage in a water-based spray as a 4-inch wide band over the row at planting time, behind the planter shoe and in front of the press wheel to achieve shallow incorporation. Use a minimum of 40 gallons of total spray volume per acre.	Do not apply more than 4.5 pints (2.25 lbs. a.i.) of Quali-Pro Chlorpyrifos 4E per acre or make more than one soil application per season. Do not use rutabaga tops for food or feed purposes. Do not make a foliar application within 10 days of a soil application. Do not apply within 30 days of harvest.

**WHEAT**

**Worker Restricted-Entry Interval:** Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours unless PPE required for early entry is worn.

For use only in Arizona, California, Colorado, Idaho, Kansas, Minnesota, Montana, Nebraska, New Mexico, Nevada, North Dakota, Oklahoma, Oregon, South Dakota, Texas, Utah, Washington, and Wyoming.

For use to control aphids (including Russian wheat aphid), wheat midge, brown wheat mite, grasshoppers, army cutworms and to provide suppression of other cutworm species.

<b>PEST</b>	<b>DOSAGE</b>	<b>TIMING/SPECIAL DIRECTIONS</b>
Aphids (including Russian wheat aphid, Greenbug, English grain aphid), Brown wheat mite, Grasshoppers	1/2 to 1 pint/acre (0.25 – 0.5 lb. a.i./acre)	From emergence to flowering, treat when 15 – 20% of tillers are infested. From flowering to early milk stage, treat when 20% or more of tillers are infested.
Wheat midge (Orange wheat blossom midge)	1 pint/acre (0.5 lb. a.i./acre)	Treatment is recommended when 75% of the wheat heads have emerged from the boot and when midge adults are found in the crop (1 midge per 4-5 heads). Application timing is critical to ensure good control. If possible, apply in the late afternoon or early evening when temperatures exceed 50° F and wind speed is less than 7 mph.
Army cutworms, other cutworm species (suppression only)	1 pint/acre (0.5 lb. a.i./acre)	Control may be reduced under high temperature conditions (greater than 80° F), under dry soil conditions, or if larvae are more than 1/2 inch long. Treat when field counts or crop injury indicates that damaging pest populations are developing or present. A second application of 1 pint/acre may be made for additional control.

Mix the required dosage with water and apply in a minimum of 2 gallons per acre finished spray volume. Apply using aerial (fixed-wing or helicopter) or power-operated ground spray equipment.

**RESTRICTIONS:** Do not make more than 2 applications per season. Do not apply within 14 days of harvest for forage and hay and within 28 days of harvest for grain and straw. Do not allow livestock to graze or otherwise feed on treated forage within 14 days of application. Do not feed straw from treated wheat within 28 days of application.

### TURF AND OTHER NONRESIDENTIAL OUTDOOR USES

Quali-Pro Chlorpyrifos 4E is an emulsifiable concentrate for control of pests located around industrial buildings (turf and ornamentals), road medians (turf and ornamentals), and golf course turf only. Pests controlled by Quali-Pro Chlorpyrifos 4E are listed in the following tables. Quali-Pro Chlorpyrifos 4E is compatible with fungicides, insecticides, and miticides commonly recommended except for alkaline materials such as Bordeaux mixtures and lime. A small jar compatibility test should always be conducted using proper proportions of chemicals and water to check for physical compatibility prior to tank mixing.

**Precautions and Restrictions:** Keep out of fish pools and other bodies of water. Do not treat vegetable gardens. Do not allow livestock to graze in treated areas. Do not feed treated grass cuttings (hay) or seed screenings to livestock or use hay for livestock bedding. Do not use Quali-Pro Chlorpyrifos 4E in poultry houses.

### ORNAMENTALS AROUND INDUSTRIAL BUILDINGS AND ROAD MEDIANS

Quali-Pro Chlorpyrifos 4E may be used to treat evergreens, vines, flowers, shrubs, shade and flowering trees, non-bearing fruit, and nut and citrus trees found around industrial buildings and road medians infested with pests listed in the following table. Dilute Quali-Pro Chlorpyrifos 4E with water according to the directions given in the table and apply using suitable hand- or power-operated spray equipment. Ensure complete and uniform coverage. Uniform coverage is critical for effective insect and mite control. Apply a coarse spray to thoroughly wet both upper and lower leaf surfaces and infested limb and trunk areas. Attempt to penetrate dense foliage, but avoid over-spraying to the point of excessive runoff. Treat when pests appear and repeat at 7- to 10-day intervals, if needed. For application timing and other specific use information, consult your State Agricultural Experiment Station or Extension Service specialist.

**Note:** Environmental factors have significant effects on phytotoxic expression. Quali-Pro Chlorpyrifos 4E has been tested on numerous ornamental plants without causing serious phytotoxicity at listed use rates. Some varieties of azaleas, camellias, poinsettias, rose bushes, or variegated ivy have shown varying degrees of phytotoxicity following treatment with Quali-Pro Chlorpyrifos 4E. Before treating large numbers of plants (especially those previously listed), treat a small block of plants and observe for 7 to 10 days to determine phytotoxic potential.

**Note:** The user assumes responsibility for determining if Quali-Pro Chlorpyrifos 4E is safe to treated plants under commercial-growing conditions.

PEST*	Amount of Quali-Pro Chlorpyrifos 4E in Water to Make		Specific Directions
	Per Acre	100 Gallons	
Adelgids: (Cooley, Eastern spruce galls, Pine bark), Ants (except fire ants, carpenter ants, harvester ants, and pharaoh ants), Aphids: (Apple, Chrysanthemum, Cottonwood, Elm leaf, Peach, Rose, Spirea, Woolly), Armyworms: (Fall, Yellowstriped), Bagworms <sup>1</sup> , Boxelder bugs, Cankerworms, Catalpa sphinx, Chiggers (for control of chiggers in golf courses, road medians, and industrial sites only), Mealy bugs, Elm spanworms, Fall webworms <sup>2</sup> , Grasshoppers, Green fruitworms, Hornworms, Jackpine budworms, Juniper webworms, Katydid, Lace bugs, Leafhoppers, Leafrollers <sup>3</sup> , Maple leafcutters <sup>4</sup> , Mites <sup>5</sup> : (Clover, Red spider, Southern red, Spruce spider, Twospotted spider), Oleander caterpillars, Orange tortrix, Periodical cicada, Plant bugs, Poplar tentmaker, Psyllids, Puss caterpillars, Rose chafers, Sawflies, exposed: (Pin oak, Pine, Redheaded), Sowbugs, Spittlebugs, Spring elm caterpillars, Springtails, Spruce budworms: (Eastern, Western), Tent caterpillars: (Eastern, Western, Forest), Thornbug, Walnut caterpillars, Whiteflies, Yellow-necked caterpillars	1 pt. – 1 qt. (0.5 - 1 lb. a.i.)	8-16 fl. oz.	<sup>1</sup> Treat when bagworm larvae are small and actively feeding. <sup>2</sup> Direct spray into web and immediately adjacent foliage for control of fall webworms. <sup>3</sup> For effective control of leafrollers, spray before leaves are tightly rolled. <sup>4</sup> Apply spray to maple leafcutter larvae as cases are being formed for effective control. Do not treat sugar maple trees intended for maple syrup production. <sup>5</sup> For effective control of spider mites when large numbers of eggs are present, apply a 2 <sup>nd</sup> spray 3-5 days in the South or 7-10 days in the North after initial treatment to control newly hatched nymphs.
Armyworms: (Beet), Beetles: (Fuller rose, Native elm bark <sup>1</sup> ), Browntail moth, Cutworms, Leafhoppers, Mahogany webworms, Mealybugs, Mimosa webworms, Moths: (Browntail, Cypress tip, Douglas fir tussock, European pine shoot, Gypsy <sup>2</sup> , Holly bud, Nantucket pine tip, Pandora, Pitch pine tip, Subtropical pine tip, Tussock, Oakworms: (California, Orangestriped, Redhumped), Redhumped caterpillars, Thrips: (Exposed), Weevils: (Blackvine <sup>3</sup> , Pine production, Yellow poplar)	1 qt. (1 lb. a.i.)	1 pt. (0.5 lb. a.i.)	<sup>1</sup> Make applications in the spring or early summer to reduce twig and branch feeding by bark beetles. <sup>2</sup> To kill migrating and invading gypsy moth larvae, treat trunks and foliage. <sup>3</sup> Blackvine weevils are night feeders. Late-afternoon spraying will maximize control.
Foliar-feeding beetles: (Blister leaf, Cottonwood leaf <sup>1</sup> , Elm leaf, Flea, Fuller rose, Japanese, June, Willow leaf)	1 qt. (1 lb. a.i.)	1 pt. (0.5 lb. a.i.)	<sup>1</sup> Use Quali-Pro Chlorpyrifos 4E in water to control cottonwood leaf beetle larvae and adults infesting cottonwoods. Make the treatment when field counts indicate damaging beetle populations are developing or present.

PEST*	Amount of Quali-Pro Chlorpyrifos 4E in Water to Make		Specific Directions
	Per Acre	100 Gallons	
Borers <sup>1</sup> , Clearwing moths: (Ash, Dogwood, Lesser peachtree, Lilac, Oak, Rhododendron), Metallic wood: (Bronze birch, Flathead appletree, Twolined chestnut), Longhorned beetles: (Locust, Red oak), Cranberry girdler larvae <sup>2</sup> , Leafminers, Needleminers: (Jeffrey pine, Lodgepole pine, Spruce), Scale insects <sup>3</sup> : (Cottonycushion, Cottony maple, Euonymus, Fletcher, Florida wax, Golden oak, Hemispherical, Lecanium, Magnolia, Oak kermes, Oak lacanium, Oystershell, Pine needle, San Jose, Tea, White birch, White peach)	1 qt. (1 lb. a.i.)	1 qt. (1 lb. a.i.)	<sup>1</sup> For borers, apply Quali-Pro Chlorpyrifos 4E to the trunks and lower limbs of trees and shrubs when the adults begin to emerge. Consult your State Agricultural Experiment Station or Extension Service specialist for proper time to treat. Apply uniformly a coarse, low-pressure spray. Pheromone traps may aid in detection of adult clearwing moths. <sup>2</sup> Apply 1 qt. of Quali-Pro Chlorpyrifos 4E for cranberry girdler larvae. Direct spray at the base of tree using 50 gallons of water per acre. Irrigate immediately after applications for soil penetration of 1-2 inches. Treat after egg laying during the summer. <sup>3</sup> Time applications for control of scale insects when crawlers or first two stages of settled nymphs are present.
Borers: (Cottonwood, Peachtree <sup>1</sup> )	1 qt. (1 lb. a.i.)	3 qts. ** (3 lbs. a.i.)	<sup>1</sup> For peachtree borers, apply Quali-Pro Chlorpyrifos 4E in water to flowering trees and shrubs of the genus Prunus as a trunk spray before newly hatched larvae enter the trees. Apply as a coarse, low-pressure spray. Thoroughly wet all bark areas from ground level to scaffold limbs. ** When using the 3-quart-per-100-gallon dilution, do not exceed 1 qt. of Quali-Pro Chlorpyrifos 4E per acre.
Beetles <sup>1</sup> : (Includes wood-infesting, Ambrosia, Anobiidae, Black turpentine, Cottonwood leaf, Elm leaf, European elm bark, Flea, Fuller rose, Japanese, June, Native elm bark <sup>2</sup> , Southern pine, Willow leaf)	1 qt. (1 lb. a.i.)	2 gals. ***	<sup>1</sup> For preventative treatment, apply the spray to the main trunk of trees in the early spring or when threat of attack exists from nearby infested trees. For remedial treatment, apply the spray to the main trunk of infested trees when damage occurs but before adult beetles begin to emerge. <sup>2</sup> To prevent native elm bark beetles from overwintering in uninfested trees, apply Quali-Pro Chlorpyrifos 4E in water to the bottom 9 ft. of the trunk. Wet the trunk thoroughly but do not spray to runoff. Care should be taken to apply the spray right to the base of the root flare. Application can be made with either a backpack mistblower or a hydraulic pressure sprayer from spring through early fall. *** When using the 2-gallon-per-100-gallon dilution, do not exceed 1 qt. of Quali-Pro Chlorpyrifos 4E per acre.

\*Superscripts refer to **Specific Directions**.



**FOR COMMERCIAL ORNAMENTALS IN NURSERIES, FOR USDA QUARANTINE USE ONLY, AND FOR DIRECT MOUND AND/OR DRENCH APPLICATION ONLY**

Quali-Pro Chlorpyrifos 4E may be used for USDA quarantine use only to treat containerized, potted, or balled and burlapped nursery stock to control the insects in the soil attached to the roots of these plants. Completely submerge the container with drain holes or root ball stabilized by burlap in a tank containing diluted Quali-Pro Chlorpyrifos 4E. Do not remove burlap wrap or plastic containers with drain holes prior to submerging. Keep the container or root ball submerged until complete soil saturation has occurred, normally about 30 seconds.

**Precautions:** During all operations (submerging, drenching, injecting), wear a chemical-resistant apron in addition to other PPE listed for applicators and other handlers. Make applications in a well-ventilated area.

**Note:** Environmental factors have significant effects on phytotoxic expression. Quali-Pro Chlorpyrifos 4E has been tested on numerous ornamental plants without causing serious phytotoxicity at listed use rates. However, because of the numerous varieties grown, treat a small group of plants at the listed rate under the anticipated growing conditions and observe for at least 7 days to determine phytotoxic potential before treating a larger number of plants.

**Note:** The professional user assumes responsibility for determining if Quali-Pro Chlorpyrifos 4E is safe to treated plants under commercial-growing conditions.

PEST*	Amount of Quali-Pro Chlorpyrifos 4E in Water to Make		Specific Directions
	1 Gallon	100 Gallons	
Fire Ants <sup>1</sup>	1/25 fl. oz.	4 fl. oz.	<sup>1</sup> As an alternative to submerging potted plants, dilute 4 oz. of Quali-Pro Chlorpyrifos 4E in 100 gallons of water. Apply this dilution to the point of runoff twice daily for 3 consecutive days. Do not remove burlap wrap or container from plants prior to treatment.
White Grubs <sup>2</sup> Weevils <sup>2</sup> (such as Blackvine)	2/3 fl. oz.	2 qts.** (2 lbs. a.i.)	<sup>2</sup> An alternative treatment to submerging containerized plants is to drench the container with the diluted insecticide solution applying approximately 10 to 12 fl. oz. of diluted insecticide solution per gallon of container size (4 to 5 fl. oz./100 cubic inches of container). The container media should be pre-moistened with irrigation or rainfall before drenching. Do not remove container from plants prior to treatment. <sup>3</sup> An alternate treatment to submerging balled and burlapped plants is to inject Quali-Pro Chlorpyrifos 4E into the root ball. Equally distribute 1 to 3 quarts of the dilute Quali-Pro Chlorpyrifos 4E solution per cubic foot of soil volume through an injection rod inserted into the soil ball surrounding the plant roots. Uniform distribution of the insecticide throughout the soil of the root ball is critical for effective control. It is recommended that the injection rod be inserted in at least 4 equally spaced locations around the stem of the plant at a 30- to 45-degree angle from the plant between the stem and the upper, outer perimeter of the ball. This technique has been shown to be most effective with small root balls (up to 1.5 ft in diameter). Larger root balls may require more injection points to ensure thorough soil distribution of the insecticide. The injection rod should be coupled to a flow meter to monitor the correct volume applied per root ball using an injection pressure of at least 30 psi. The application should be made such that splash-back and runoff are minimized.
Coffee root mealybug <sup>2</sup>	1/6 fl. oz.	1 pt. (0.5 lb. a.i.)	<sup>2</sup> Do not exceed more than 1 quart (1 lb. a.i.) of Quali-Pro Chlorpyrifos 4E per acre.

\*Superscripts refer to **Specific Directions**.

**ORNAMENTALS IN INDUSTRIAL PLANT SITES AND ROAD MEDIANS  
(DORMANT SPRAY OF TREE PESTS)**

Quali-Pro Chlorpyrifos 4E may be used as a dormant or delayed-dormant spray at the rates indicated to control the listed insects. Quali-Pro Chlorpyrifos 4E may be used without oil; however, oil is recommended to control additional pests such as the European red mite.

For high-volume (dilute) sprays (200 to 600 gallons of spray mixture per acre), tank mix the specified dosage with 1 to 2 gallons of a petroleum spray oil recommended for dormant use in 100 gallons of water. Spray the entire tree to runoff using suitable ground spray equipment.

For low-volume (concentrate) sprays (less than 200 gallons of spray mixture per acre), use the same amount of Quali-Pro Chlorpyrifos 4E as for a dilute spray and apply in a manner that will ensure thorough coverage of the trees. Use oil as recommended by your State Agricultural Experiment Station or Extension Service specialist.

**Precautions:** Do not apply until rain or irrigation have replenished soil moisture such that bark and twigs are not desiccated since cold, dry conditions may cause Quali-Pro Chlorpyrifos 4E plus oil to infuse trees, resulting in bud damage or drop.

**Restrictions:** Make only one application during the dormant season except for the control of the apple ermine moth. Do not allow meat or dairy animals to graze in treated areas.

PEST*	Amount of Quali-Pro Chlorpyrifos 4E in Water to Make			Specific Directions
	1 Gallon	3 Gallons	100 Gallons	
Aphids: (Mealy plum, Rosy apple, Woolly apple), Borers: (Peach twig), Cutworms: (Climbing), Leafrollers: (Pandemis), Pear psylla adults, Plant bugs, Scale: (San Jose)	1/12 – 1/6 fl. oz.	1/4 - 1/2 fl. oz.	1/2 - 1 pt.	Tank mix with 1 to 2 gallons of a petroleum spray oil recommended for dormant use in 100 gallons of water.
Apple ermine moth	1/12 fl. oz.	1/4 fl. oz.	1/2 pt.	For control of <i>Malus</i> species, make 2 applications at a 7- to 14-day interval in combination with a petroleum spray oil at the rate of 2% to 4% (v/v) in a spray to wet application to ensure thorough coverage of all stems and branches. When using tank mixtures, follow all label directions for the mixing partner (oil). Use appropriate application equipment and spray volumes to ensure complete coverage of the plant(s), or control will be compromised.

\*Superscripts refer to **Specific Directions**.

**TURF, INDUSTRIAL PLANT, AND ROAD MEDIAN OUTDOOR USES**

Apply Quali-Pro Chlorpyrifos 4E to control the pests listed in the following table at the listed dosages and in accordance with the directions given below or as recommended by your local Agricultural Extension Service specialist. Dilute Quali-Pro Chlorpyrifos 4E in water and apply as a coarse, low-pressure spray using suitable application equipment. Except as noted, thoroughly water immediately after treatment to wash the insecticide into the turf. The area to be treated should be moist at the time of treatment. Spray when pests first appear; re-treat when needed.

PEST*	Amount of Quali-Pro Chlorpyrifos 4E per		Specific Directions
	1000 sq. feet	Acre	
Ticks <sup>1</sup> : (American dog, Cattle fever, Gulf coast, Lone star) (for control of ticks in golf courses, road medians, and industrial plant sites only)	¼ fl. oz.	1 ½ pts.	<sup>1</sup> For control of ticks, treat soil and other areas likely to serve as harborage sites for ticks that have removed themselves from their host. Spray surfaces to be treated until wet, but do not create excessive runoff. <b>Note:</b> This application is intended as a premise spray only. Do not use as a direct spray on livestock or any sites that may come in contact with livestock.
Ants (except fire ants, carpenter ants, harvester ants, and pharaoh ants), Armyworms: (Beet, Fall, Yellowstriped), Chiggers <sup>1</sup> (for control of chiggers in golf courses, road medians, and industrial plant sites only), Chinch bugs, Clover mites, Cutworms, Crickets, Deer ticks <sup>2</sup> (for control of ticks in golf courses, road medians, and industrial sites only), Earwigs, Fiery skipper, Gnats, Grasshoppers, Greenbug aphids, June beetles, Leafhoppers, Lucerne moths, Millipedes, Mites: (Clover, Bermuda-grass stunt, Formula grass, Winter grain), Pillbugs, Sod webworms <sup>3</sup> : (Lawn moths), Sowbugs, Ticks <sup>1</sup> (for control of ticks in golf courses, road medians, and industrial plant sites only)	¾ fl. oz.	1 qt.	<sup>1</sup> Apply Quali-Pro Chlorpyrifos 4E for area control of ticks and chiggers infesting golf course turf, turf in road medians, and industrial plant sites where these pests are present and create a nuisance or a possible public health problem. Do not allow public use of treated areas during application or until spray has dried. Apply Quali-Pro Chlorpyrifos 4E in water at the rate of ½ pint/acre (equivalent to 1/6 fl. oz. per 1000 sq. ft.) using a hydraulic sprayer, mist applicator, knapsack sprayer, or other suitable hand- or power-operated spray equipment. Treat low underbrush, grassy areas, weeds, and ground surface and debris using enough spray volume to obtain thorough coverage, usually 40-100 gals./acre. <sup>2</sup> Apply Quali-Pro Chlorpyrifos 4E in water at the rate of 1 quart per acre or ¾ fl. oz. per 1000 sq. ft. for control of deer ticks. Treat low underbrush, turf, grassy areas, weeds, and ground surface and debris using enough spray volume to obtain thorough coverage. <sup>3</sup> For sod webworms, watering or mowing of the treated area should be delayed for 24 hours after treatment.
European crane fly	1 fl. oz.	1 qt.	
Turfgrass weevil ( <i>Hyperodes</i> ) <sup>1</sup>	1 ½ fl. oz.	1 qt.	<sup>1</sup> Make application to problem areas in mid-April and again in mid-May or as recommended by your local Agricultural Extension Service specialist.
White grubs <sup>1</sup> : (Black turfgrass atenianus, European chafer, Japanese beetle larvae, Southern and Northern masked chafer)	1 ½ - 3 fl. oz.	1 qt.	<sup>1</sup> Spray when white grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by your local Agricultural Extension Service specialist. Immediately after spraying, irrigate the treated area with ½ to 1 inch of water to wash the insecticide deep into the thatch or into the underlying soil.
Billbug adults such as (Bluegrass, Denver, Hunting)	¾ - 3 fl. oz.	1 qt.	Spray early in the season when adult billbugs first appear.
Mole crickets	1 ½ fl. oz.	1 qt.	For mole crickets in golf course turf, turf in road medians, and industrial plant site turfgrass, apply through high-pressure injection or other suitable subsurface placement application equipment. Depending on the application equipment used, follow the manufacturer's recommendation for calibration and the volume of spray per acre needed to provide control or as recommended by your local Agricultural Extension Service specialist. Apply when young nymphs are active.

\*Superscripts refer to **Specific Directions**.

**OUTSIDE SURFACES AND AROUND INDUSTRIAL PLANT SITES (SUCH AS AROUND WAREHOUSES, FOOD PROCESSING, AND FOOD MANUFACTURING SITES)**

Quali-Pro Chlorpyrifos 4E may be applied as a residual spray to and around outside surfaces of nonresidential buildings and structures. Permitted areas of use include fences, pre-construction foundations, refuse dumps, outside of walls, and other areas where pests congregate or have been seen. Do not allow adults, children, or pets to contact treated surfaces until sprays have dried. Keep out of fish pools and other bodies of water. Do not feed treated grass cuttings (hay) or seed screenings to livestock or use treated hay for livestock bedding. Do not treat vegetable gardens. Repeat treatment as needed to maintain effectiveness. Unless prohibited by a product's label, users, at their own discretion, can tank mix pesticides currently labeled for similar use patterns. Always perform a small jar compatibility test using proper proportions to check for physical compatibility prior to tank mixing. Do not tank mix this product with products containing dichlorvos (DDVP).

PEST*	Amount of Quali-Pro Chlorpyrifos 4E in Water to Make			Specific Directions
	1 Gallon	10 Gallons	50 Gallons	
	For Band Treatment <sup>†</sup>			
Ants (except fire ants, carpenter ants, harvester ants, and pharaoh ants), Beetles, Boxelder bugs (for other true bugs), Clover mites, Crickets, Earwigs, Elm leaf beetles (adults), Firebrats, Millipedes, Pillbugs, Silverfish, Sowbugs, Spiders (excluding black widow and brown recluse spiders), Springtails, Ticks (for control of ticks in golf courses, road medians, and industrial plant sites only)	¼ tsp.	¾ fl. oz.	4 fl. oz.	<sup>†</sup> To help prevent infestation of non-residential buildings, treat a band of soil 6 to 10 ft. wide around and adjacent to buildings including the building foundation to a height of 2 to 3 ft. where pests are active and may find entrance. Use 4 fl. oz. of Quali-Pro Chlorpyrifos 4E per 50 gals. of water and apply as a coarse spray at the rate of about 10 gals. of spray mixture per 1000 sq. ft. to thoroughly and uniformly wet the band area.
<b>For Outside Surfaces</b>				
	1 1/3 fl. oz.	13 1/3 fl. oz.	2 qts.**	**Do not exceed 1 qt. of Quali-Pro Chlorpyrifos 4E per acre.

\*Superscripts refer to **Specific Directions**.

**ANTS (EXCEPT FIRE ANTS, CARPENTER ANTS, HARVESTER ANTS, AND PHARAOH ANTS) AND MISCELLANEOUS PESTS**

PEST	Amount of Quali-Pro Chlorpyrifos 4E in Water to Make			Specific Directions
	1 Gallon	3 Gallons	100 Gallons	
Ants (except fire ants, carpenter ants, harvester ants, and pharaoh ants), Cockroaches (American, Asian, Brownbanded, German, Oriental, Smokey brown, Wood), Sowbugs, Springtails	1/6 fl. oz.	½ fl. oz.	1 pt.	

# STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in original container in secured dry storage area. Prevent cross-contamination with other pesticides and fertilizers. Do not store above 100° F for extended periods of time. Storage below 20° F may result in formation of crystals. If product crystallizes, store at 50° F to 70° F and agitate to redissolve crystals. If container is damaged or spill occurs, use product immediately or dispose of product and damaged container as indicated below.

**PESTICIDE DISPOSAL:** Open dumping is prohibited. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your state pesticide or environmental control agency or the hazardous waste representative at the nearest EPA Regional Office for guidance.

## **CONTAINER HANDLING:**

**Nonrefillable Container (five gallons or less):** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

## **LIMITATION OF WARRANTY AND LIABILITY**

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following **CONDITIONS, DISCLAIMER OF WARRANTIES** and **LIMITATIONS OF LIABILITY**.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Makhteshim Agan of North America, Inc. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of Makhteshim Agan of North America, Inc. is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Makhteshim Agan of North America, Inc. disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at Makhteshim Agan of North America, Inc.'s election, the replacement of product.

Quali-Pro is a registered trademark of Makhteshim Agan of North America, Inc.